

Number **2**

2016

Volume 49

49 (1) 57–97

ISSN 1337-9984 (Print)

ISSN 1338-0095 (Online)

Slovak Journal of **Animal Science**



NATIONAL AGRICULTURAL
AND FOOD CENTRE

RESEARCH INSTITUTE FOR ANIMAL
PRODUCTION NITRA

Editorial office

Editor-in-chief:

Ladislav Hetényi, NPPC – Research Institute for Animal Production Nitra, Slovak Republic

Executive editor:

Ludmila Hanuliaková, NPPC – Research Institute for Animal Production Nitra, Slovak Republic

Technical editor:

Marta Vargová, NPPC – Research Institute for Animal Production Nitra, Slovak Republic

Editorial board

Daniel Bíro, Slovak University of Agriculture Nitra, Slovakia

Zsuzsanna Bősze, Agricultural Biotechnology Center, Gödöllő, Hungary

Jan Brouček, NPPC – Research Institute for Animal Production Nitra, Slovakia

Jozef Bulla, Slovak University of Agriculture Nitra, Slovakia

Ondrej Debrecéni, Slovak University of Agriculture Nitra, Slovakia

Andrzej Filistowicz, The Faculty of Biology and Animal Science, University of Environmental and Life Science, Wrocław, Poland

Roland Grossmann, Institute of Animal Science Mariensee, Germany

Jaroslav Olav Horbańczuk, Institute of Genetics and Animal Breeding of the Polish Academy of Sciences in Jastrzębiec n/Warsaw, Poland

Peter Chreněk, NPPC – Research Institute for Animal Production Nitra, Slovakia

Jozef Laurinčík, Constantine the Philosopher University Nitra, Slovakia

Juraj Koppel, Institute of Animal Physiology SAS, Košice, Slovakia

Alexander Vladimírovič Makarevič, NPPC – Research Institute for Animal Production Nitra, Slovakia

Peter Massanyi, Slovak University of Agriculture Nitra, Slovakia

Gábor Mészáros, University of Natural Resources and Life Sciences, Division of Livestock Sciences, Vienna, Austria

Štefan Mihina, NPPC – Research Institute for Animal Production Nitra, Slovakia

Shoukhart M. Mitalipov, Oregon Health & Science University, Beaverton, U.S.A.

Jaana Peippo, MTT Agrifood Research Finland, Jokioinen, Finland

Dana Peškovičová, NPPC – Research Institute for Animal Production Nitra, Slovakia

Juraj Pivko, NPPC – Research Institute for Animal Production Nitra, Slovakia

Josef Příbýl, Research Institute for Animal Production, Praha – Uhřetěves, Czech Republic

Ján Rafay, NPPC – Research Institute for Animal Production Nitra, Slovakia

Alexander Sirotkin, NPPC - Research Institute for Animal Production Nitra, Slovakia

Pavel Suchý, Department of Nutrition, Dietetics, Zoohygiene and Plant Products, University of Veterinary and Pharmaceutical Sciences Brno, Czech Republic

Milan Šimko, Slovak University of Agriculture Nitra, Slovakia

Peter Šútovský, University of Missouri – Columbia, U.S.A.

Vladimír Tančin, NPPC – Research Institute for Animal Production Nitra, Slovakia

Manuel Tena-Sempere, University of Córdoba, Spain

Miroslav Valent, West Virginia University, Morgantown, U.S.A.

Igor Valocký, University of Veterinary Medicine, Košice, Slovakia

Scientific English: **Shubhadeep Roychoudhury, Alexander V. Makarevič**

AIMS AND SCOPE

Slovak Journal of Animal Science (ISSN 1337-9984) is an international scientific journal that publishes original scientific papers, reviews, short communications, chronicles of important jubilees, reports of participation in important international conferences on animal science in English language.

Topic of the journal are problems of animal production, mainly in the sphere of genetics, breeding, nutrition and feeding, physiological processes of digestion, conservation and treatment of feeds, biotechnology, reproduction, ethology, ecologization of breeding, biology and breeding technology in farm animals, quality of meat, milk, wool, economy of breeding and production of farm animals, mainly: cattle, pigs, sheep, goats, horses, poultry, small farm animals and farm game. There are published also articles from the sphere of biochemistry, genetics, embryology, applied mathematical statistics as well as economy of animal production. There can be published also articles from the sphere of veterinary medicine concerning the themes of the journal.

No part of this publication may be reproduced, stored or transmitted in any form without prior written permission of the Publisher.

SJAS is published quarterly.

Online version of the journal (ISSN 1338-0095) is at disposal at <http://www.vuzv.sk/index.php/slovak-journal-of-animal-science>.
Permission of the Publisher is required to store or use electronically any material contained in this journal.

This journal is comprised in: AGRIS/FAO database (the full texts, too); CAB Abstracts; Knovel.

Slovak Journal of Animal Science is published under the authorization and direction of the National Agricultural and Food Centre - Research Institute for Animal Production Nitra, Slovak Republic.

Editorial office, orders, subscription and distribution: NPPC - RIAP Nitra, Hlohovecká 2, 951 41 Lužianky, Slovak Republic. Phone +421 37 6546 249; E-mail: editor@vuzv.sk; <http://www.vuzv.sk>; www.nppc.sk; www.sjas.sk

Filed at the Ministry of Culture of the Slovak Republic: EV 3659/09.

© NPPC - RIAP Nitra, 2016.

FERTILIZING ABILITY OF PINZGAU BULL SPERM *IN VITRO* AFTER CRYOSTORAGE

L. OLEXIKOVÁ^{1*}, E. ŠPALEKOVÁ¹, E. KUBOVIČOVÁ¹, A. V. MAKAREVICH¹, P. CHRENEK^{1,2}

¹NPPC – Research Institute for Animal Production Nitra, Slovak Republic

²Faculty of Biotechnology and Food Science, Slovak University of Agriculture in Nitra, Slovak Republic

ABSTRACT

The aim of the study was to evaluate fertilizing ability of Pinzgau bull sperm after different periods of cryostorage. The sperm samples from 16 bulls were arranged into the three groups according to cryostorage period as follows: less than 7 years (group 1), 7 to 13 years (group 2) and 14 or more years (group 3). Straws with frozen sperm doses were thawed in a water bath and the motile fraction of sperm was selected by modified swim-up method. Motile spermatozoa ($2 \times 10^6 \cdot \text{ml}^{-1}$) were co-incubated with matured bovine oocytes in fertilization drops (IVF-TALP medium) under mineral oil at 39 °C in presence of 25 µg/ml heparin for 20 hours. Totally, 739 oocytes were used in the *in vitro* fertilization test. Following fertilization the presumptive zygotes were stained with DAPI nuclear stain and the status of chromatin was examined under fluorescence microscope. Penetration of the sperm head into the ooplasm of oocyte and fertilization (formation of pronuclei) were evaluated. No significant difference in fertilizing ability among the experimental groups was found. The rates of penetrated plus fertilized eggs for group 1, 2 and 3 were 79.31, 76.40 and 76.14 %, respectively. Significantly higher pronuclear formation rate was observed in the group 1, where 72.9 % of eggs had two visible pronuclei at 20 hours following incubation with sperm compared to 65.72 and 47.21 % in groups 2 and 3, respectively. Also, 4 % of fertilized eggs from the group 1 reached syngamy of both pronuclei, probably due to a faster formation of pronuclei, whilst no syngamy was observed in groups 2 and 3. In conclusion, period of cryostorage had no direct influence on penetrating and fertilizing ability of Pinzgau bull sperm doses. However, it is assumed that cryostorage length can affect the speed of pronucleus formation.

Key words: Pinzgau; bull; sperm; fertilizing ability; cryopreservation

INTRODUCTION

Cryostorage of bull semen is widely used as a useful tool to improve reproduction. In dairy cattle, the majority of inseminations are done with frozen-thawed semen. Cryopreserved semen plays a major role in genetic improvement, economics of breeding programs in the livestock industry, and preservation of endangered species and breeds. Sperm cryopreservation methods are the most efficient way for storing animal genetic resources. It is well known, that cryopreservation is stressful for spermatozoa. Although the purpose of sperm cryopreservation is to preserve sperm function and fertility, the freezing-thawing process inevitably

causes their damage and reducing fertility. During freezing and thawing spermatozoa are subjected to various stresses such as cold shock, osmotic and oxidative stress and intracellular ice crystal formation. These cause disruption of sperm integrity. Particularly, membrane structure alterations (Krogenaes *et al.*, 1994; Januskauskas *et al.*, 2003), and sperm function alterations after thawing (Garner *et al.*, 1997; Thomas *et al.*, 1998; Pons-Rejraji *et al.*, 2009) were described. In addition, excessive mitochondrial activity induces the generation of reactive oxygen species (Chatterjee and Gagnon, 2001) that affects functions of cellular compounds and organelle (Bilodeau *et al.*, 2000).

Some evidences also exist about cryodamages

*Correspondence: E-mail: olexikova@vuzv.sk
Lenka Olexiková, NPPC – Research Institute for Animal Production Nitra
Hlohovecká 2, 951 41 Lužianky, Slovak Republic
Tel.: +421 37 6546 196

Received: January 13, 2016
Accepted: May 13, 2016

to sperm DNA integrity after cryopreservation (Gandini *et al.*, 2006; Waterhouse *et al.*, 2010). These damages ultimately result in reduced sperm motility, viability and fertilizing ability (Watson, 2000). It is assumed, that once frozen semen in the liquid nitrogen can be potentially stored unlimited time. But only scarce information was published about the influence of long-term cryostorage on quality of frozen-thawed bull semen. In humans, it was proved that sperm can be useful after even more than 28 years of cryostorage (Clarke *et al.*, 2006). But some alterations may be related to long period of storage. For example the percentage of motile sperm was lower after long-term (9–13 years) compared to short-term (1–5 years) storage (Edelstein *et al.*, 2008).

The aim of the study was to evaluate fertilizing ability of Pinzgau bull sperm after different periods of cryostorage in liquid nitrogen.

MATERIAL AND METHODS

Procedure of *in vitro* fertilization (IVF)

Bovine ovaries were collected at a local abattoir and transported to the laboratory within 3 hours at 25–30 °C. Ovaries were rinsed with 70 % ethanol and a sterile saline solution. Cumulus-oocyte complexes (COCs) were acquired by follicular fluid aspiration from visible large follicles (2–10 mm in diameter) on the surface of ovaries. Collected COCs with pale homogeny cytoplasm and complete cumulus were selected and washed twice in maturation medium E199 with HEPES (Biowest). For maturation, E199 medium with glutaMAX (Gibco), sodium pyruvate (0.25 mmol.l⁻¹), gentamycin (0.05 mg.ml⁻¹), fetal bovine serum 10 % (FBS, BioWhittaker, Verviers, Belgium) and FSH/LH (1/1 I.U., Pluset, Lab. Calier, Barcelona, Spain) was used. Oocytes were matured at 39 °C in a humidified atmosphere with 5 % CO₂ in air for 24 h. Matured oocytes were partially cleaned from cumulus cells by vortexing during 3 min. Frozen-thawed insemination doses from Pinzgau bulls were used for *in vitro* fertilization of oocytes. Insemination doses from 16 bulls of a proven fertility were used in experiments. Straws were thawed in a water bath at 36 °C and the motile fraction of sperm was separated in Sperm-TALP medium using modified swim-up procedure. The sperm fraction was then resuspended in IVF-TALP medium and chosen sperm concentration was adjusted. Following maturation, the presumptive matured oocytes were placed into 100 µl drops of fertilization medium IVF-TALP under sterile mineral oil and incubated with 2×10^6 .ml⁻¹ spermatozoa in the presence of 25 µg/ml heparin (Sigma-Aldrich, Germany) at 39 °C in a humidified atmosphere with 5 % CO₂ in air for 20 hours.

Following this time, the presumptive zygotes

were cleaned off from the excessive sperm and remaining cumulus cells by vortexing, washed twice in Sperm-TALP medium and in the PBS with 0.6 % polyvinylpyrrolidone. Zygotes were immediately fixed in 4 % formalin during 10 min and then covered with a drop of Vectashield anti-fade medium containing DAPI stain (chromatin staining; Vector Laboratories, Burlingame, CA, USA) and mounted between coverslip and microslide and stored at 4 °C until fluorescence analysis. Stained embryos were checked under a Leica fluorescence microscope using specific filter with wavelength for blue fluorescence and x20 magnification objective. Penetration of sperm head into the ooplasm of oocyte and formation of pronuclei (fertilizing ability) were evaluated.

Statistical analysis

Differences in distribution of oocytes into either the penetrated, fertilized (presence of pronuclei) or non-penetrated among the groups of different cryostorage period were analyzed by Pearson's Chi-square test of independence. All calculations were performed using the SAS software package (SAS Institute, 2001).

RESULTS AND DISCUSSION

Fertilizing ability estimated by *in vitro* fertilization test may be a good tool to predict potential fertility of a bull (Larsson and Rodríguez-Martínez, 2000). *In vitro* fertilization test was previously used by several authors to test fertilizing ability of sperm as a homologous system with cow oocytes and bull sperm (Fazeli *et al.*, 1997; Zhang *et al.*, 1998; Marquant-Le Guienne *et al.*, 1990) or as heterologous system with sperm of different species using cow oocytes. Such heterologous system was, for example, used to test ram (García-Alvarez *et al.*, 2009; Makarevich *et al.*, 2011), antelope (Roth *et al.*, 1998), stallion (Bromfield *et al.*, 2013) and dolphin (Sánchez-Calabuig *et al.*, 2015) sperm fertilizing ability. Major advantage of these *in vitro* tests is that the testing is not so expensive and labor intensive procedure as testing by artificial insemination. The *in vitro* fertilization technique can be very accurate to assess the sperm fertility, because the procedure evaluates the spermatozoa-oocyte interactions occurring during fertilization process.

In our previous study (Makarevich *et al.*, 2011) we used co-incubation of ram spermatozoa with bovine oocytes with intact *zona pellucida* in the *in vitro* fertilization test, where about 53–55 % of the oocytes were penetrated by ram semen and about 30–33 % of them were fertilized. Similarly García-Alvarez *et al.* (2009) documented that such heterologous *in vitro* fertilization test can be useful to predict the *in vivo* fertility of rams.

Fertilizing ability is very individual property of bull (Ax and Lent, 1987; Merckies *et al.*, 2000).

To minimize individual effect of bull and underline the effect of time we created experimental groups from minimum three bulls, according to the length of cryostorage period.

No significant difference in fertilizing ability (rate of penetrated and fertilized eggs) among the three experimental groups was found. The total rates of penetrated and fertilized eggs for group 1, 2 and 3 were 79.31; 76.40 and 76.14 %, respectively (Table 1). Similarly, period of bull sperm storage had no influence on the occurrence of polyspermy after *in vitro* fertilization. According to period of storage 6.3 %, 6.5 % and 5.1 % of fertilized eggs were polyspermic in groups 1, 2 and 3, respectively (Table 1). Similar rate of polyspermy was reported by Santos with co-workers (2008), where the polyspermy rates varied from 4.1 to 11.1 %. It was proved, that polyspermy after bovine *in vitro* fertilization is mainly influenced by individuality of a bull, heparin dose used (Marquant-Le Guenne *et al.*, 1990) and also related to oocyte quality (Santos *et al.*, 2008).

On the other hand, significant differences were found in the rate of pronuclei formation. Significantly

higher pronuclear formation was observed in the group 1 (stored for 1–6 years), where 72.9 % of eggs had two visible pronuclei at 20 hour after sperm addition compared to 65.72 and 47.21 % in groups 2 and 3, respectively (Table 1). Also, syngamy of both pronuclei was observed in 4 % of fertilized eggs in the group 1, whilst no syngamy was observed in groups 2 and 3. This difference may be probably due to faster pronuclear formation in the group 1.

Slower decondensation of sperm heads occurring in groups 2 and 3 may be related to chromatin structure alteration, described in cryostored human (Royere *et al.*, 1991) and boar (Hamamah *et al.*, 1990) spermatozoa. In particular, Royere with co-workers (1991) concluded that freezing-thawing procedure may alter the DNA/nuclear protein relationships and impair the fertilizing ability of human sperm. Similarly, Hamamah *et al.* (1990), basing on their results, hypothesize that sperm chromatin after freezing-thawing may be overcondensed; this overcondensation may be associated with the lower conception rates obtained using human and porcine semen after cryostorage.

Table 1: Fertilizing ability of Pinzgau bull sperm after cryostorage

Groups	Period of storage (years)	No. bulls	No. oocytes	Penetrated and fertilized (N) %	Non-penetrated (N) %	Oocytes with two pronuclei (N) %	Polyspermic oocytes from fertilized (n/N) %
1	1 to 6	3	203	(161) 79.3	(42) 20.7	(148) 72.9	(10/158) 6.3
2	7 to 13	7	339	(259) 76.4	(80) 23.6	(227) 67.0	(16/243) 6.5
3	14 to 18	6	197	(150) 76.1	(47) 23.9	(93) 47.2	(5/98) 5.1
total		16	739	(570) 77.1	(169) 22.9	(468) 63.3	(31/499) 6.2

CONCLUSION

In conclusion, length of cryostorage had no direct influence on *in vitro* penetrating and fertilizing ability of cryopreserved Pinzgau bull sperm. However, long-term cryostorage might affect speed of pronucleus formation and syngamy in *in vitro* fertilized eggs.

ACKNOWLEDGEMENTS

This work was supported by the Slovak Research and Development Agency from the grant APVV-14-0043 and from the Ministry of Agriculture of the Slovak republic: RPVV3 “Výskumné postupy pre zachovanie

biodiverzity”. We also thank to Slovak Biological Services Inc., Lužianky for providing with Pinzgau bull insemination doses.

REFERENCES

- AX, R.-L. – LENT, R. W. 1987. Glycosaminoglycans as probes to monitor differences in fertility of bulls. *Journal of Dairy Science*, vol. 70, 1987, p. 1477–1486.
- BILODEAU, J. F. – CHATTERJEE, S. – SIRARD, M. A. – GAGNON, C. 2000. Levels of antioxidant defenses are decreased in bovine spermatozoa after a cycle of freezing and thawing. *Molecular Reproduction and Development*, vol. 55, no. 3, 2000, p. 282–288.

- BROMFIELD, E. G. – AITKEN, R. J. – GIBB, Z. – LAMBOURNE, S. R. – NIXON, B. 2013. Capacitation in the presence of methyl- β -cyclodextrin results in enhanced zona pellucida-binding ability of stallion spermatozoa. *Reproduction*, vol. 147, no. 2, 2013, p. 153–166.
- CLARKE, G. N. – LIU, D. Y. – BAKER, H. W. G. 2006. Recovery of human sperm motility and ability to interact with the human zona pellucida after more than 28 years of storage in liquid nitrogen. *Fertility and Sterility*, vol. 86, 2006, p. 721–722.
- EDELSTEIN, A. – YAVETZ, H. – KLEIMAN, S. E. – HAUSER, R. – BOTCHAN, A. – PAZ, G. – YOGEV, L. 2008. Effect of long-term storage on deoxyribonucleic acid damage and motility of sperm bank donor specimens. *Fertility and Sterility*, vol. 90, 2008, p. 1327–1330.
- FAZELI, A. R. – ZHANG, B. R. – STEENWEG, W. – LARSSON, B. – BEVERS, M. M. – VAN DEN BROEK, J. – RODRIGUEZ-MARTINEZ, H. – COLENBRANDER, B. 1997. Relationship between sperm-zona pellucida binding assays and the 56-day nonreturn rate of cattle inseminated with frozen-thawed bull semen. *Theriogenology*, vol. 48, no. 5, 1997, p. 853–863.
- GANDINI, L. – LOMBARDO, F. – LENZI, A. – SPANO, M. – DONDERO, F. 2006. Cryopreservation and sperm DNA integrity. *Cell and Tissue Banking*, vol. 7, 2006, p. 91–98.
- GARCÍA-ALVAREZ, O. – MAROTO-MORALES, A. – MARTINEZ-PASTOR, F. – FERNANDEZ-SANTOS, M. R. – ESTESO, M. C. – PEREZ-GUZMAN, M. D. – SOLER, A. J. 2009. Heterologous *in vitro* fertilization is a good procedure to assess the fertility of thawed ram spermatozoa. *Theriogenology*, vol. 71, 2009, p. 643–650.
- GARNER, D. L. – THOMAS, C. A. – JOERG, H. W. – DEJARNETTE, J. M. – MARSHALL, C. E. 1997. Fluorometric assessments of mitochondrial function and viability in cryopreserved bovine spermatozoa. *Biology of Reproduction*, vol. 57, no. 6, 1997, p. 1401–1406.
- HAMAMAH, S. – ROYÈRE, D. – NICOLLE, J. C. – PAQUIGNON, M. – LANSAC, J. 1990. Effects of freezing-thawing on the spermatozoon nucleus: a comparative chromatin cytophotometric study in the porcine and human species. *Reproduction Nutrition and Development*, vol. 30, no. 1, 1990, p. 59–64.
- CHATTERJEE, S. – GAGNON, C. 2001. Production of reactive oxygen species by spermatozoa undergoing cooling, freezing, and thawing. *Molecular Reproduction and Development*, vol. 59, no. 4, 2001, p. 451–458.
- JANUSKAUSKAS, A. – JOHANNISSON, A. – RODRIGUEZ-MARTINEZ, H. 2003. Subtle membrane changes in cryopreserved bull semen in relation with sperm viability, chromatin structure, and field fertility. *Theriogenology*, vol. 60, no. 4, 2003, p. 743–758.
- KROGENAES, A. – ANDERSEN BERG, K. – HAFNE, A. L. – ENGELAND, E. 1994. Membrane alterations in bull spermatozoa after freezing and thawing and after *in vitro* fertilization. *Acta Veterinaria Scandinavica*, vol. 35, no. 1, 1994, p. 17–26.
- LARSSON, B. – RODRÍGUEZ-MARTÍNEZ, H. 2000. Can we use *in vitro* fertilization tests to predict semen fertility? *Animal Reproduction Science*, vol. 60–61, 2000, p. 327–336.
- MAKAREVICH, A. V. – SPALEKOVA, E. – OLEXIKOVA, L. – LUKAC, N. – KUBOVICOVA, E. – HEGEDUSOVA, Z. 2011. Functional characteristics of ram cooling-stored spermatozoa under the influence of epidermal growth factor. *General Physiology and Biophysics*, vol. 30, Special Issue, 2011, S36–S43.
- MARQUANT-LE GUIENNE, B. – HUMBLLOT, P. – THIBIER, M. – THIBAUT, C. 1990. Evaluation of bull semen fertility by homologous *in vitro* fertilization tests. *Reproduction Nutrition and Development*, vol. 30, no. 2, 1990, p. 259–266.
- MERKIES, K. – LARSSON, B. – KJELLÉN, L. – ZHANG, B. R. – BUHR, M. M. – RODRÍGUEZ-MARTÍNEZ, H. 2000. Relationship between heparin binding to spermatozoa and the fertility of dairy bulls. *Theriogenology*, vol. 54, no. 8, 2000, p. 1249–1258.
- PONS-REJRAJI, H. – BAILEY, J. L. – LECLERC, P. 2009. Cryopreservation affects bovine sperm intracellular parameters associated with capacitation and acrosome exocytosis. *Reproduction, Fertility and Development*, vol. 21, no. 4, 2009, p. 525–537.
- ROTH, T. L. – WEISS, R. B. – BUFF, J. L. – BUSH, L. M. – WILDT, D. E. – BUSH, M. 1998. Heterologous *in vitro* fertilization and sperm capacitation in an endangered African antelope, the scimitar-horned oryx (*Oryx dammah*). *Biology of Reproduction*, vol. 58, no. 2, 1998, p. 475–482.
- ROYERE, D. – HAMAMAH, S. – NICOLLE, J. C. – LANSAC, J. 1991. Chromatin alterations induced by freeze-thawing influence the fertilizing ability of human sperm. *International Journal of Andrology*, vol. 14, no. 5, 1991, p. 328–332.
- SÁNCHEZ-CALABUIG, M. J. – DE LA FUENTE, J. – LAGUNA-BARRAZA, R. – BELTRÁN-BREÑA, P. – MARTÍNEZ-NEVADO, E. – JOHNSTON, S. D. – RIZOS, D. – GUTIÉRREZ-ADÁN, A. – PÉREZ-GUTIÉRREZ, J. F. 2015. Heterologous murine and bovine IVF using bottlenose dolphin (*Tursiops truncatus*) spermatozoa. *Theriogenology*, vol. 84, no. 6, 2015, p. 983–994.
- SANTOS, P. – CHAVEIRO, A. – SIMÕES, N. – MOREIRA DA SILVA, F. 2008. Bovine oocyte

- quality in relation to ultrastructural characteristics of zona pellucida, polyspermic penetration and developmental competence. *Reproduction in Domestic Animals*, vol. 43, no. 6, 2008, p. 685–689.
- THOMAS, C. A. – GARNER, D. L. – DEJARNETTE, J. M. – MARSHALL, C. E. 1998. Effect of cryopreservation of bovine sperm organelle function and viability as determined by flow cytometry. *Biology of Reproduction*, vol. 58, no. 3, 1998, p. 786–793.
- WATERHOUSE, K. E. – GJELDNE, A. – TVERDAL, A. – DE ANGELIS, P. M. – FARSTAD, W. – HÅRD, M. – KOMMISRU, E. 2010. Alterations of sperm DNA integrity during cryopreservation procedure and *in vitro* incubation of bull semen. *Animal Reproduction Science*, vol. 117, no. 1-2, 2010, p. 34–42.
- WATSON, P. F. 2000. The causes of reduced fertility with cryopreserved semen. *Animal Reproduction Science*, vol. 60-61, 2000, p. 481–492.
- ZHANG, B. R. – LARSSON, B. – LUNDEHEIM, N. – RODRIGUEZ-MARTINEZ, H. 1998. Sperm characteristics and zona pellucida binding in relation to field fertility of frozen-thawed semen from dairy AI bulls. *International Journal of Andrology*, vol. 21, 1998, p. 207–216.

CRYOPRESERVATION OF AMNIOTIC FLUID STEM CELLS DERIVED FROM ZOBOR RABBIT

M. KOVÁČ^{1,2*}, B. KULÍKOVÁ^{1,3}, J. VAŠÍČEK^{1,2}, P. CHRENEK^{1,2}

¹Faculty of Biotechnology and Food Science, Slovak University of Agriculture in Nitra, Slovak Republic

²NPPC – Research Institute for Animal Production Nitra, Slovak Republic

³Department of Zoology and Anthropology, Constantine the Philosopher University, Nitra, Slovak Republic

ABSTRACT

The aims of our preliminary study were to evaluate the viability and to investigate the occurrence of early and late apoptosis of fresh, frozen-thawed and thawed-cultured rabbit amniotic fluid stem cells (AFSCs). Rabbit AFSCs were isolated from young (4–6 months old) Zobor rabbit females ($n=3$), cultured in endothelial basal medium (EBM-2) to passage 3 (P3) and frozen. One month after cryopreservation AFSCs were thawed and cultured for 72 hours. We identified the apoptotic (Annexin-V - AnV; Yo-Pro-1 - YoP) and dead (propidium iodide - PI) AFSCs using flow cytometry and evaluated the viability of fresh (F-P3), frozen-thawed (F-T) and thawed-cultured (72 hours of post-thaw *in vitro* culture – T-72h) AFSCs. To examine the phenotype of P3 AFSCs we performed an antibody-based staining of surface markers CD29, CD44 and CD45. We have found decreased ($P < 0.05$) viability in F-T cells (77.5 ± 2.2 % and 75.2 ± 0.7 % for AnV/PI⁺ and YoP/PI⁺, respectively), compared to F-P3 (92.5 ± 2.2 and 89.45 ± 1.3 for AnV/PI⁺ and YoP/PI⁺, respectively). However, viability of T-72h cells was similar to F-P3 (90.9 ± 1.5 % and 84.6 ± 1.7 % for AnV/PI⁺ and YoP/PI⁺, respectively). Our AFSCs were both CD44⁺ (95.98 ± 1.51 %) and CD29⁺ (92.33 ± 4.46) positive but CD45⁺ (0.73 ± 0.55 %) negative, according to defined MSCs phenotype.

Key words: rabbit; amniotic fluid; stem cells; viability; cryopreservation

INTRODUCTION

Stem cells are the focus of great interest because of their potential for therapy in a wide variety of conditions. Conditions possibly treatable by stem-cell transplants include genetic defects, tissue and organ replacement, autoimmune disease and malignancies (Dzidosz *et al.*, 2016). These cells with high differentiation capacities are able to give rise to tissue derivatives of three germ layers; ecto-, meso- and endoderm lineages. They can be spontaneously differentiated through the formation of embryoid bodies (Valli *et al.*, 2010), and can also undergo targeted differentiation under appropriate culture conditions (De Coppi *et al.*, 2007a).

Amniotic fluid collected routinely via amniocentesis allows us to bank stem cells. These AFSCs can be proliferated in culture as needed and

offer a convenient source for autologous transplantations (Abdulrazzak *et al.*, 2013). They can be isolated from mid-term amniotic fluid (AF) of both human and rodents and cultured up to 300 passages (De Coppi *et al.*, 2007b; Phermathai *et al.*, 2010). Many authors declare that human AFSC are cultured easily, remain viable over many passages and tolerate cryopreservation very well (Da Sacco *et al.*, 2011; De Coppi *et al.*, 2007a; Young *et al.*, 2015; Chen *et al.*, 2014). Besides human, they have been isolated from various species including dogs, pigs, horses and buffaloes (Chen *et al.*, 2011; Choi *et al.*, 2013; Dev *et al.*, 2013; Filioli Uranio *et al.*, 2011; Iacono *et al.*, 2012; Yadav *et al.*, 2011).

Stem cells are characterized by the presence of surface markers that are associated with self-renewal without differentiation. The flow cytometry has been a primary technique for the identification of stem cells

***Correspondence:** E-mail: 3mkovac3@gmail.com
 Michal Kováč, Institute of Farm Animal Genetics and Reproduction
 NPPC – Research Institute for Animal Production Nitra
 Hlohovecká 2, 951 41 Lužianky, Slovak Republic
 Tel.: +421 37 6546 285 Fax: +421 37 6546 189

Received: March 15, 2016

Accepted: May 23, 2016

by detection of these markers (Dziadosz *et al.*, 2016). Previously described AFSCs also exhibit intermediate characteristics between embryonic and adult stem cells. They are devoid of ethical controversies and are safer in clinical applications (Chen *et al.*, 2011; De Coppi *et al.*, 2007b; Pozzobon *et al.*, 2010). In the past, scientists classified AFSCs as pluripotent, according to the marker expression. However, the inability to form tumors upon transplantation positioned AFSCs as broad multipotent (Canazi and De Coppi, 2012) or potentially pluripotent stem cells (Gao *et al.*, 2013). Moreover, primary cultures of amniotic fluid contain a heterogeneous population of cells due to the direct contact of the fluid and the fetus (Cremer *et al.*, 1981; Ferdaos and Nordin, 2012; Joo *et al.*, 2012). Paebst *et al.* (2014) showed method for selection of specific cells prior to the cultivation by sorting of CD44⁺ cells from amniotic fluid.

The use of preclinical models is promising for future therapeutic possibilities (Dziadosz *et al.*, 2016). For example, Klein *et al.* (2011) transplanted AFSCs to fetal lambs and reported more rapid healing time of skin wounds. Sun *et al.* (2015) differentiated human AFSCs into keratinocytes that improved epidermal regeneration of intentionally excised wounds in mouse model. De Koninck *et al.* (2015) observed improved lung density and function in rabbit model with congenital diaphragmatic hernia after intratracheal injection of human AFSCs.

The aims of our preliminary study were to evaluate the viability and to investigate the occurrence of early and late apoptosis of fresh, frozen-thawed and thawed-cultured rabbit AFSCs.

MATERIAL AND METHODS

Cells isolation and culture

Amniotic fluid was obtained from humanely sacrificed Zbor rabbits ($n = 3$) at 23rd day of gravidity using pipette under sterile conditions and mixed with culture medium at the 5:6 ratio. The medium was composed of EBM-2 basal medium (CC-3156, Lonza, USA) supplemented with 20 % fetal calf serum (FCS), EGM-2 SingleQuots™ Kit (CC-4176, Lonza, USA) and 1 % of penicillin and streptomycin (Life Technologies, Slovakia). Five ml of the mixture was used for seeding the cells into T25 culture flasks. Cells were cultured and medium was changed at 5th day, after the cells formed adherent colonies. Subsequently, the culture medium was changed every 2nd day. Cells were dissociated after the colonies reached 80-90 % confluency using Accutase (Invitrogen, Carlsbad, CA, USA) for 5 min at 37 °C and 5 % CO₂. AFSCs were counted and seeded into new culture flasks at density of $4-5 \times 10^4$ cells/cm².

Subsequently, the cells were cultured and passaged after 3-4 days, after they reached 80-90 % confluency.

Freezing

Rabbit AFSCs were frozen in freezing solution composed of medium used for culture and 10 % DMSO (D2650, Sigma-Aldrich). 2×10^6 cells were mixed with 1.5 ml of freezing solution and placed into cryovials. Cryovials were frozen at slow rate by reducing the temperature by 1 °C per minute in Mr. Frosty container (Thermo Scientific Nalgene, Rochester, NY, USA) and stored for 24 hours at -80 °C. After 24 hours the cryovials were immersed into liquid nitrogen and stored for 1 month. Thawing of the AFSCs was rapid and performed by swirling the cryovials in water bath at 37 °C for 30-60 sec. The suspension was then slowly instilled into pre-warmed culture medium to prevent osmotic shock.

Fluorescent staining

The viability of the cells was evaluated prior to cryopreservation (F-P3), immediately after thawing (F-T) and after additional culture for 72 h (T-72 h). Annexin-V (Roche Slovakia, Slovak Republic) and Yo-Pro-1 (Molecular Probes, Lucerne, Switzerland) staining was used for detection of early and late apoptosis, respectively. The dead cells were identified by propidium iodide (Molecular Probes, Eugene, Oregon, USA) staining. At least $0.5-1 \times 10^6$ cells were washed in AnV buffer and PBS without magnesium/calcium (Life Technologies, Slovakia) for AnV and YoP staining, respectively. Washed cells were placed in 50 µl of prepared staining solution and incubated in the dark at room temperature for 20 min. At least 1×10^4 cells in each sample were analyzed by flow cytometer FACS Calibur (BD Biosciences, USA).

Surface marker detection

Flow cytometry immunostaining was performed to identify the expression of the cell surface markers CD44, CD29 and CD45. Primary antibodies were as follows: anti-rabbit CD29 (mouse IgG1, clone P4G11, Abcam, United Kingdom), anti-rabbit CD44 (mouse IgG1, clone W4/86, AbD Serotec, Germany), anti-rabbit CD45 (mouse IgG1, clone L12/201, AbD Serotec, Germany). Secondary antibodies were as follows: anti-IgG1-PE, mouse (rat IgG1, clone X-56, Miltenyi Biotec, Germany) for CD44 and CD29; anti-IgG1-APC, mouse (rat IgG1, clone X-56, Miltenyi Biotec, Germany) for CD45. Briefly, F-P3 cells were dissociated, washed with PBS and centrifuged for 5 min at $587 \times g$ at 4 °C. The pellet was resuspended in 50 µl of rabbit serum to block the Fc receptors. Cells were incubated with isotype controls, primary and secondary antibodies for 15 min on ice. At least 0.5×10^5 cells were analyzed for each CD marker.

Statistical Analysis

The statistical analysis was performed with One-Way ANOVA (Tukey-Test) for comparison of mean values $\pm$ SDs using SigmaPlot software (Systat Software Inc., Germany).

RESULTS AND DISCUSSION

Our present research was focused on isolation of rabbit AFSCs that could serve as a tool for preserving the animal genetic resources in gene bank.

Immunophenotyping of rabbit AFSCs in our study revealed that they express surface molecular markers CD44 and CD29 that are commonly used to profile MSCs. The expression of CD44 was 95.98 ± 1.51 %, which fulfil criteria of the International Society for Cellular Therapy for positive MSCs marker expression (Dominici *et al.*, 2006). The expression of CD29 was slightly lower (92.33 ± 4.46 %). Nevertheless, such positivity was considered as sufficient for accepting these cells as CD29⁺ according to other studies, which declared AFSCs as CD29 positive, without quantification (You *et al.*, 2009; Paebst *et al.*, 2014; Filioli Uranio *et al.*, 2011).

Moreover, AFSCs in our study were more positive to CD44 marker than AFSCs isolated by Slamečka and Chrenek (2013). We suggest that more than the two-fold higher proportion of cells expressing CD44 (95.98 ± 1.51 % vs 47.8 %) might be attributed to antibodies specificity. We have used anti-rabbit antibodies, whereas Slamečka and Chrenek (2013) used antibodies designed for human AFSCs. It is necessary to say, that the markers required for defining a MSC were mainly determined for human

cells, and the lack of specific antibodies for different animal species led to a great variability of approaches and results for molecular characterization of animal MSCs (Iacono *et al.*, 2015). Furthermore, hematopoietic lineage marker CD45 was used as a negative control. The cells fulfilled the criteria of Dominici *et al.* (2006) for negative marker expression (0.73 ± 0.55 %).

Viability of AFSCs was evaluated by flow cytometry fluorescent staining. AnV/PI⁻ and YoP/PI⁻ cells were gated to evaluate the overall viability. We have found decreased ($P < 0.05$) viability in F-T cells (77.5 ± 2.2 % and 75.2 ± 0.7 % for AnV and YoP, respectively), compared to F-P3 control (92.5 ± 2.2 and 89.45 ± 1.3 for AnV and YoP respectively). However, viability of T-72 h cells was similar to fresh passage (90.9 ± 1.5 % and 84.6 ± 1.7 % for AnV/PI⁻ and YoP/PI⁻, respectively).

Our results showed that freezing of stem cells increased the apoptotic and dead cell proportion. It has been reported that hematopoietic and other stem cells, particularly human embryonic stem cells, undergo apoptotic transformation during the cryopreservation process (Sangeetha *et al.*, 2010; Stroh *et al.*, 2002). MSCs also lose their viability very soon post-thaw, what can most likely be attributed to the rapid development of apoptotic processes (Berz and Colvin, 2012). Our F-T AFSCs showed more than two-fold increase in apoptotic or PI⁺ cells compared to F-P3 AFSCs (9.8 ± 1.8 % vs 3.7 ± 1.2 %; 10.5 ± 4.8 % vs 4.0 ± 1.8 % for AnV/PI⁻ and YoP/PI⁻, respectively). Pal *et al.* (2008) investigated the post-thaw behavior of MSCs and reported a rapid decrease in viability from > 80 % at 2 hours to < 40 % at 8 hours, while the cells were maintained at 4 °C.




Fig. 1: Phase contrast images of the post-thaw cultured rabbit AFSCs (Zeiss Primo Vert Inverted Microscope). 1A – endothelial-like shape (10x magnification) 1B – mesenchymal-like shape (20x magnification).

Table 1: Detection of Annexin-V, Yo-Pro-1, and propidium iodide positive/negative cells expressed in %

Amniotic fluid stem cells	AnV-/PI-	AnV+/PI-	AnV-/PI+	AnV+/PI+	YoP-/PI-	YoP+/PI-	YoP-/PI+	YoP+/PI+
Fresh	92.5 ± 2.2 ^a	0.6 ± 0.3	4.1 ± 1.7 ^a	3.7 ± 1.2 ^a	89.45 ± 1.3 ^a	2.6 ± 0.9	4.6 ± 1.7	4.0 ± 1.8 ^a
Thawed	77.5 ± 2.2 ^b	2.9 ± 0.9	8.5 ± 0.9 ^b	9.8 ± 1.8 ^b	75.2 ± 0.7 ^b	3.8 ± 1.2	6.2 ± 0.8	10.5 ± 4.8 ^b
Thawed 72 h	90.9 ± 1.5 ^a	1.2 ± 1.4	4.9 ± 1.9 ^{ab}	4.0 ± 1.9 ^a	84.6 ± 1.7 ^a	3.6 ± 0.3	3.9 ± 1.0	5.9 ± 2.4 ^{ab}

Results are expressed as means ± SEM; a vs b within the same column were statistically significant at P < 0.05.

The decrease in viability may be associated also with DMSO addition. It is a small amphipathic molecule that penetrates via plasma membrane of cells and acts as a strong hydrogen bond disrupter and hence exerts colligative effects (Ruiz-Delgado *et al.*, 2009). It affects multiple organ systems with wide spectrum of toxicities. However, a DMSO concentration of 10 % in stem cell concentrate is still considered to be standard in most centers around the world. Interestingly, the maximal possible cryopreservation time span is still unknown (Berz and Colvin, 2012).

CONCLUSION

Although the freezing of rabbit AFSCs negatively affects the viability and lead to increase in apoptotic processes, our AFSC were able of post-thaw proliferation. Moreover, after additional 72 hours of culture the proportion of dead and apoptotic cells was reduced. The AFSCs morphology did not change post-thaw and following culture. These preliminary results could potentially serve for cryopreservation of these cells with subsequent use for gene bank purposes as genetic resources or for tissue engineering.




Fig. 2: Proportion of apoptotic, dead and viable AFSCs - P3 fresh (F-P3); frozen-thawed (F-T); thawed-cultured for 72 hours (T-72 h). 2A – proportion of living AnV-/PI- and YoP-/PI- AFSCs. 2B – Necrotic and apoptotic AnV+/PI+ and YoP+/PI+ AFSCs.

ACKNOWLEDGEMENT

This work was supported by the grants APVV-14-0043 and VEGA 1/0611/15 coordinated by the Slovak Research and Development Agency.

REFERENCES

- ABDULRAZZAK, H. – DE COPPI, P. – GUILLOT, P. V. 2013. Therapeutic potential of amniotic fluid stem cells. *Current Stem Cell Research & Therapy*, vol. 8 (2), 2013, p. 117–124.
- BERZ, D. – COLVIN, G. 2012. Cryopreservation of hematopoietic and non-hematopoietic stem cells – A Review for clinician, New advances in stem cell transplantation. 1st ed., Rijeka: Intech, 2012, 582 p., ISBN 978-953-51-0013-3.
- CANAZI, M. – DE COPPI, P. 2012. CD117 (+) amniotic fluid stem cells: state of the art and future perspectives. *Organogenesis*, vol. 8 (3), 2012, p. 77–88.
- CHEN, J. – LU, Z. – CHENG, D. – PENG, S. – WANG, H. 2011. Isolation and characterization of porcine amniotic fluid-derived multipotent stem cells. *PLoS One*, vol. 6 (5), 2011, p. 19964.
- CHEN, Z. – CHAN, M. K. – STELCHENKO, N. – WANG, F. – LIU, L. – PERLE, M. A. – BASCH, R. S. – YOUNG, B. K. 2014. Heterogeneity of stem cells in human amniotic fluid. *Journal of Regenerative Medicine*, vol. 3 (1), 2014, p. 1–8.
- CHOI, S. – CHOI, H. – KIM, K. J. – LEE, D. – LEE, J. – PARK, J. – KIM, E. – LI, X. – OH, H. – LEE, D. – KIM, M. K. 2013. Isolation of canine mesenchymal stem cells from amniotic fluid and differentiation into hepatocyte-like cells. *In Vitro Cellular & Developmental Biology - Animal*, vol. 49 (1), 2013, p. 42–51.
- CREMER, M. – TREISS, I. – CREMER, T. – HAGER, D. – FRANKE, W. W. 1981. Characterization of cells of amniotic fluids by immunological identification of intermediate-sized filaments: presence of cells of different tissue origin. *Human Genetics*, vol. 59 (4), 1981, p. 373–379.
- DA SACCO, S. – DE FILIPPO, R. E. – PERIN, L. 2011. Amniotic fluid as a source of pluripotent and multipotent stem cells for organ regeneration. *Current Opinion in Organ Transplantation*, vol. 16 (1), 2011, p. 101–105.
- DE COPPI, P. – BARTSCH, G. – SIDDIQUI, M. M. – XU, T. – SANTOS, C. C. – PERIN, L. – MOSTOSLAVOVSKY, G. – SERRE, A. C. – SNYDER, E. Y. – YOO, J. J. – FURT, M. E. – SOKER, S. – ATALA, A. 2007a. Isolation of amniotic fluid stem cell lines with potential for therapy. *Nature Biotechnology*, vol. 25 (1), 2007a, p. 100–106.
- DE COPPI, P. – CALLEGARI, A. – CHIAVEGATO, A. – GESPAROTTP, L. – PICCOLI, M. – TAIANI, J. – POZZOBON, M. – BOLDRIN, L. – OKABE, M. – COZZI, E. – ATALA, A. – GAMBA, P. – SARTORE, S. 2007b. Amniotic fluid and bone marrow derived mesenchymal stem cells can be converted to smooth muscle cells in the cryo-injured rat bladder and prevent compensatory hypertrophy of surviving smooth muscle cells. *Journal of Urology*, vol. 177 (1), 2007b, p. 369–376.
- DE KONINCK, P. – TOELAN, J. – ROUBLIOVA, X. – CARTER, S. – POZZOBON, M. – RUSSO, F. M. – RICHTER, J. – VANDERSLOTEN, P. J. – VERBEKEN, E. – DE COPPI, P. – DEPREST, J. 2015. The use of human amniotic fluid stem cells as an adjunct to promote pulmonary development in a rabbit model congenital diaphragmatic hernia. *Prenatal Diagnostics*, vol. 35 (9), 2015, p. 833–840.
- DEV, K. – GIRI, S. K. – KUMAR, A. – YADAV, A. – SINGH, B. – GAUTAM, S. K. 2012. Derivation, characterization and differentiation of buffalo (*Bubalus bubalis*) amniotic fluid derived stem cells. *Reproduction in Domestic Animals*, vol. 47 (5), 2013, p. 704–711.
- DOMINICI, M. – LE BLANC, K. – MUELLER, I. – SLAPER-CORTENBACH, I. – MARINI, F. – KRAUSE, D. – DEANS, R. – KEATING, A. – PROCKOP, D. J. – HORWITZ, E. 2006. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy*, vol. 8 (4), 2006, p. 315–317.
- DZIADOSZ, M. – BASCH, R. S. – YOUNG, B. K. 2016. Human amniotic fluid: a source of stem cells for possible therapeutic use. *American Journal of Obstetrics & Gynecology*, vol. 214 (3), 2016, p. 321–327.
- FERDAOS, N. – NORDIN, N. 2012. Human amniotic fluid cells and their future perspectives. *Regenerative Research*, vol. 1 (2), 2012, p. 14–19.
- FILIOLI URANIO, M. – VALENTINI, L. – LANGE-CONSIGLIO, A. – CAIRA, M. – GUARICCI, A. – L'ABBATE, A. – CATACHIO, C. R. – VENTURA, M. – CREMONESI, F. – DELL'AQUILA, M. E. 2011. Isolation, proliferation, cytogenetic, and molecular characterization and *in vitro* differentiation potency of canine stem cells from foetal adnexa: a comparative study of amniotic fluid, amnion, and umbilical cord matrix. *Molecular Reproduction and Development*, vol. 78 (5), 2011, p. 361–373.
- GAO, L. – THILAKAVATHY, K. – NORDIN, N. 2013. A plethora of human pluripotent stem cells. *Cell Biology International*, vol. 37 (9), 2013, p. 875–887.

- IACONO, E. – BRUNORI, L. – PIRRONE, A. – PAGLIARO, P. P. – RICCI, F. – TAZZARI, P. L. – MERLO, B. 2012. Isolation, characterization and differentiation of mesenchymal stem cells from amniotic fluid, umbilical cord blood and Wharton's jelly in the horse. *Reproduction*, vol. 143 (4), 2012, p. 455–468.
- IACONO, E. – ROSSI, B. – MERLO, B. 2015. Stem cells from Foetal Adnexa and Fluid in Domestic Animals: An update in their features and clinical application. *Reproduction in Domestic Animals*, vol. 50 (3), 2015, p. 353–364.
- JOO, S. – KO, I. K. – ATALA, A. – YOO, J. J. – LEE, S. J. 2012. Amniotic fluid-derived stem cells in regenerative medicine research. *Archives of Pharmacal Research*, vol. 35 (2), 2012, p. 271–280.
- KLEIN, J. D. – TURNER, C. G. – STEIGMAN, S. A. – AHMED, A. – ZURAKOWSKI, D. – ERIKSSON, E. – FAUZA, D. O. 2011. Amniotic mesenchymal stem cells enhance normal fetal wound healing. *Stem Cells and Development*, vol. 20 (6), 2011, p. 969–976.
- PAEBST, F. – PIEHLER, D. – BREHM, W. – HELLER, S. – SCHROECK, C. – TARNOK, A. – BURK, J. 2014. Comparative immunophenotyping of equine multipotent mesenchymal stromal cells: an approach toward a standardized definition. *Cytometry A*, vol. 85 (8), 2014, p. 678–687.
- PAL, R. – HANWATE, M. – TOTTEY, S. M. 2008. Effect of holding time, temperature and different parenteral solutions on viability and functionality of adult bone marrow-derived mesenchymal stem cells before transplantation. *Journal of Tissue Engineering and Regenerative Medicine*, vol. 2 (7), 2008, p. 436–444.
- PERMATHAI, T. – ODGLUN, Y. – JULAVIJITPHONG, S. – TITAPANT, V. – CHUENWATTANA, P. – VATANASIRI, C. – PATTANAPANYASAT, K. 2010. A novel method to derive amniotic fluid stem cells for therapeutic purposes. *BMC Cell Biology*, vol. 11, 2010, p. 79.
- POZZOBON, M. – GHIONZOLI, M. – DE COPPI, P. 2010. ES, iPS, MSC, and AFS cells. Stem cells exploitation for Pediatric Surgery: current research and perspective. *Pediatric Surgery International*, vol. 26 (1), 2010, p. 3–10.
- RUIZ-DELGADO, G. J. – MANCÍAS-GUARRA, C. – TAMEZ-GÓMEZ, E. L. – RODRÍGUEZ-ROMO, L. N. – LÓPEZ-OTERO, A. – HERNÁNDEZ-ARIZPE – GÓMEZ-ALMAGUER, D. – RUIZ-ARGÜELLES, G. J. 2009. Dimethyl sulfoxide-induced toxicity in cord blood stem cell transplantation: report of three cases and review of the literature. *Acta Haematologica*, vol. 122 (1), 2009, p. 1–5.
- SANGEETHA, V. M. – KALE, V. P. – LIMAYE, L. S. 2010. Expansion of cord blood CD34 cells in presence of zVADfmk and zLLYfmk improved their *in vitro* functionality and *in vivo* engraftment in NOD/SCID mouse. *PloS ONE*, vol. 5 (8), 2010.
- SLAMEČKA, J. Jr. – CHRENEK, P. 2013. Rabbit amniotic fluid as a potential alternative source of broadly multipotent stem cells. *Slovak Journal of Animal Science*, vol. 46, no. 1, p. 11–15.
- STROH, C. – CASSENS, U. – SAMRAJ, A. K. – SIBROWSKI, W. – SCHULZE-OSTHOFF, K. – LOS, M. 2002. The role of caspases in cryoinjury: caspase inhibition strongly improves the recovery of cryopreserved hematopoietic and other cells. *The FASEB Journal*, vol. 16 (12), 2002, p. 1651–1653.
- SUN, Q. – LI, F. – LI, H. – CHEN, R. H. – GU, Y. Z. – CHEN, Y. – LIANG, H. S. – YOU, X. R. – DING, S. S. – GAO, L. – WANG, Y. L. – QIN, M. D. – ZAHNG, X. G. 2015. Amniotic fluid stem cells provide considerable advantages in epidermal regeneration: B7H4 creates a moderate inflammation microenvironment to promote wound repair. *Scientific Reports*, vol. 23 (5), 2015, p. 11560.
- VALLI, A. – ROSNER, M. – FUCHS, C. – SIEGEL, N. – BISHOP, C. E. – DOLZNIG, H. MÄDEL, U. – FEICHTINGER, W. – ATALA, A. – HENGSTSCHLÄGER, M. 2010. Embryoid body formation of human amniotic fluid stem cells depends on mTOR. *Oncogene*, vol. 29 (7), p. 966–977.
- YADAV, P. S. – MANN, A. – SINGH, V. – YASHVEER, S. – SHARMA, R. K. – SINGH, I. 2011. Expression of pluripotency genes in buffalo (*Bubalus bubalis*) amniotic fluid. *Reproduction in Domestic Animals*, vol. 46 (4), 2011, p. 705–711.
- YOU, Q. – TONG, X. – GUAN, Y. – ZHANG, D. – HUANG, M. – ZHANG, Y. – ZHENG, J. 2009. The biological characteristics of human third trimester amniotic fluid stem cells. *Journal of International Medical Research*, vol. 37 (1), 2009, p. 150–112.
- YOUNG, B. K. – CHAN, M. K. – LIU, L. – BASCH, R. S. 2015. Amniotic fluid as a source of multipotent cells for clinical use. *Journal of Perinatal Medicine*, vol. 44 (3), 2016, p. 333–337.

COMPARATIVE STUDY BETWEEN ISA BROWN AND FULANI ECOTYPE CHICKENS SUPPLEMENTED WITH HUMIC ACID

A. O. LALA^{1*}, N. OKWELUM¹, K. O. BELLO¹, N. A. FAMAKINDE¹, M. O. ALAMU²

¹Institute of Food Security, Agricultural Research and Environmental Resources, Federal University of Agriculture, Abeokuta, Ogun State, Nigeria

²Department of Animal Nutrition, Federal University of Agriculture, Abeokuta, Ogun State, Nigeria

ABSTRACT

A 32 week study was carried out to determine the influence of humic acid supplementation on the growth, production and egg qualities of ISA Brown and Fulani ecotype chickens. One hundred and eighty day old chicks of 90 each of ISA Brown and Fulani ecotype were randomly distributed in a 2x3 factorial arrangement of these breeds (ISA Brown and Fulani ecotype chickens) and three treatment groups 1) control with no humic acid supplementation, 2) supplementation with 1 mL.L⁻¹ humic acid and 3) 2 mL.L⁻¹ humic acid in drinking water. The six groups were replicated thrice with 10 chicks in each replicate. Humic acid supplementation ($p < 0.05$) increased body weight, improved feed conversion ratio, reduced mortality, increased hen-day production and egg weight in both ISA Brown and Fulani ecotype chickens (FEC). Feed intake ($p < 0.05$) increased in ISA Brown chickens with humic acid supplementation, while it decreased in Fulani ecotype chickens at eight weeks of age. At 20 weeks of age, both breeds consumed ($p < 0.05$) less feed with humic acid supplementation. pH of the different segments of gastrointestinal tract was ($p < 0.05$) lowered by humic acid supplementation. Shell thickness was significantly ($p < 0.05$) increased at 2 mL.L⁻¹ humic acid supplementation level in ISA Brown and FEC with ISA Brown groups having ($p < 0.05$) thicker shells when compared with FEC. High density lipoprotein ($p < 0.05$) increased across the treatment groups in ISA Brown, while these values ($p < 0.05$) decreased in ISA Brown groups supplemented with humic acid. Low density lipoprotein ($p < 0.05$) increased with humic acid supplementation in ISA Brown and FEC. The results of the present study indicated that the use of humic acid at 1 mL.L⁻¹ improved the body weight gain, overall feed intake and hen-day production in ISA Brown and Fulani ecotype chickens. However, better nutrient utilization was observed in Fulani ecotype chickens when compared with ISA Brown chickens.

Key words: chicks; growth; laying hens; lipid profile; egg qualities; humic acid

INTRODUCTION

The poultry industry has experienced tremendous growth; this growth has been with exotic chickens. The indigenous chicken genetic resources in Nigeria have been seriously endangered owing to genetic erosion through the rapid replacement by exotic breeds. The local chicken constitutes about 80 percent of the 120 million poultry birds found in Nigeria (FMA&RD, 2006). These chickens are also known for their adaptation superiority in terms of their resistance to endemic diseases and other harsh environmental conditions

(Horst, 1989). The Nigerian indigenous chickens are thought to be suitable for the development of layer strains for the tropical environment since they possess some inherent advantages which include good fertility and hatchability, flavour, colour and texture of meat and egg that is preferred by local consumers, high degree of adaptability to prevailing conditions, high genetic variance in their performance, hardiness, disease tolerance, ease of rearing and ability to breed naturally (Adebambo *et al.*, 2009).

Fulani ecotype chicken (FEC) is native to the Fulani tribe in the middle belt and northern parts of Nigeria.

***Correspondence:** E-mail: funmilala02@yahoo.com
 Adebukunola Olufunmilayo Lala, Institute of Food Security, Agricultural
 Research and Environmental Resources, Federal University of Agriculture,
 P. M. B. 2240 Abeokuta, Ogun State, Nigeria
 Tel.: +2348060395255 Fax: +23439244299

Received: November 23, 2015

Accepted: March 2, 2016

They are known to be superior in live weight than any other chicken ecotype within Nigeria (Olawunmi *et al.*, 2008). The indigenous chickens provide immense benefits for keepers but their productivity is significantly hindered by genetic and management problems. The Nigerian Fulani Ecotype (FEC) chicken has been reported to have great potential for genetic improvement in growth and reproductive performance (Fayeye *et al.*, 2005). The need to improve on the productivity of FEC chickens could be achieved through the use of growth promoters. Improving productivity of this chicken ecotype through growth promoter will improve the economy of the local poultry farmers.

There has been increased interest in alternative natural growth promoters due to microbial resistance and residual effect of antibiotics, one of which is organic acid (Kopecky *et al.*, 2012; Sheikh *et al.*, 2011).

Organic acids are non-ionised, weak acid which can penetrate the bacterial cell wall and disrupt the normal physiology of some types of bacteria (Dhawale, 2005). Humic acid is one of the main components of humic substances which include humus, humic acid, fulvic acid, ulmic acid and trace minerals and it is the most well known of the group (Yildiz *et al.*, 2006). The mode of their action is related to the reduction of pH in the upper intestinal tract, interfering with the growth of undesirable bacteria and modifying the intestinal flora (Kirchgesner and Roth, 1982). Organic acid has been reported to have the beneficial effects of improving feed conversion ratio, growth performance, enhancing mineral absorption (Kral *et al.*, 2011; Galik and Rolinec, 2011; Petruska *et al.*, 2012). Humic substances have many beneficial effects like antibacterial, antiviral and anti-inflammatory effects. They also improve immune system, reduce odour in faeces, cause a reduction in stress and play a role in liver function (Islam *et al.*, 2005). The use of humic substances in animal brings a number of advantages for animal health and productive performance (Eren *et al.*, 2000). According to FAO (2000), there is still a considerable and largely unexploited potential for increased production from local birds through improved management. The aim of this study was to determine the effects of humic acid on performance, egg traits and egg lipid profile of ISA Brown and Fulani ecotype chickens.

MATERIAL AND METHODS

Test supplement: Humic acid used is a product of Dynapharmlab Associate SDN. BHD. It contains chelated micronutrients, nitrogen 2.35 %, phosphorus 4.44 %, potassium 1.75 %, magnesium 0.36 %, iron 867 ppm, manganese 223 ppm, copper 144 ppm, zinc

153 ppm, boron 0.011 %, molybdenum 0.002 % and humic acid 0.68 %.

Experimental management and design

A total of 180 one day-old female chicks of 90 ISA Brown and 90 Fulani ecotype chicks were randomly allocated into 3 treatment groups: 1) control with no humic acid supplementation, 2) supplementation with 1 mL.L⁻¹ humic acid and 3) 2 mL.L⁻¹ humic acid supplementation in drinking water. Each treatment group was replicated thrice with 10 chicks per replicate. The experiment was conducted for 32 weeks period with all the chickens kept under uniform management conditions throughout the experimental period. Diets were formulated for the chick (0-8 weeks), grower (8-20 weeks) and layer phases of the experiment. Nutrient compositions of diets were determined according to the AOAC (1990) as shown in Table 1. Drinking water was replaced every day.

Growth performance: Birds were weighed per replicate at the beginning and on weekly basis. Feed intake was recorded weekly and calculated as g per bird; mortality was recorded as it occurred.

Gastrointestinal tract pH measurement

At the end of 8 weeks, two hens from each replicate were randomly selected and slaughtered by cervical dislocation. The gastrointestinal tract (GIT) was removed; 10 g each of contents from the crop, gizzard, duodenum, jejunum and ileum were collected in sterilized bottles (1:10 dilution) and the value of the pH for the different segments of the GIT was measured immediately using a digital pH meter.

Production performance: Between 28th and 32nd week of the experiment, eggs were collected daily and egg production was calculated on a hen-day basis. Eggs were weighed. Feed intake was recorded weekly and calculated as g per hen per day.

Egg quality measurement

A total of 18 eggs per treatment, 6 eggs per replicate were randomly selected on weekly basis within 24 hours of lay. The eggs were individually weighed on an electronic balance, the length and width was measured using Vernier calliper to determine the egg shape index (ESI). The eggs were broken into a flat surface where yolks were separated from the whites and then weighed. The shells were carefully washed of any adhering albumen, air dried and weighed. The thickness of each shell was determined using a micrometer screw gauge. Yolk and shell percentages were determined in relation to the egg weight. Yolk colour was determined by matching with one of the matching bands of the Roche colour fan of 12 graded colours. Albumen height

Table 1: Composition of experimental basal diet (%)

Ingredient	Starter (0 – 8 weeks)	Grower (9 – 20 weeks)	Layers
Maize	53.00	58.00	47.00
Groundnut cake	12.00	9.00	5.00
Soyabean meal	18.00	8.00	20.00
Fish meal	4.00	0.00	1.00
Wheat offal	6.90	19.00	15.80
Bone meal	3.00	3.00	2.50
Oyster shell	2.00	2.00	8.00
*Premix	0.25	0.25	0.25
Salt	0.30	0.25	0.25
Lysine	0.30	0.25	0.10
Methionine	0.25	0.25	0.10
Total	100.00	100.00	100.00
Crude protein (%)	22.80	16.93	17.06
Metabolizable energy (MJ.kg ⁻¹)	11.72	11.11	10.93

*Premix for chick per kg of diet: Vitamin A 10,000 iu; vitamin D3 900 iu; copper 0.1 mg; vitamin E 50.0 mg; manganese 8.5 mg; vitamin K 2.0 mg; iron 75.0 mg; vitamin B1 2.0 mg; folic acid 5.0 mg; vitamin C 26.0 mg; pantothenic acid 20.0 mg; vitamin B6 2.0 mg; choline 1200 mg; vitamin B12 0.01 mg; niacin 50 mg; zinc 70 mg; biotin 0.2 mg.

*Premix for grower per kg of diet: Vitamin A 8,000 iu; vitamin D3 1,200 iu; copper 2.0 mg; vitamin E 31.0 mg; manganese 80 mg; vitamin B2 10.0 mg; pantothenic acid 150.0 mg; iodine 1.2 mg; selenium 0.1 mg; cobalt 2 mg.

*Premix for layer per kg of diet: Vitamin A 10,000 iu; vitamin D3 200 iu; vitamin E 100 iu; vitamin K 20 mg; thiamine 15 mg; riboflavin B2 40 mg; pyridoxine B6 15 mg; niacin 150 mg; pantothenic acid 50 mg; folic acid 5 mg; biotin 0.2 mg; choline chloride 12 mg; antioxidant 1.25 g; manganese 0.8 g; zinc 0.5 g; iron 0.2 g; copper 0.5 g; iodine 0.12 g; selenium 2 mg; cobalt 2 mg.

was measured using tripod spherometer and Haugh unit was calculated according to Haugh (1937).

Yolk lipid profile: At the end of the 32nd week of the experiment, 3 eggs from each replicate were randomly chosen to determine yolk lipid profile. The eggs were hard-cooked, allowed to cool, after which the weight of the boiled eggs were noted. The yolks were carefully separated, weighed and crumbled. 1 g sample of each yolk was homogenized with 15 ml of chloroform-methanol 2:1 (v/v), thoroughly mixed and filtered. Egg homogenate filtrates were designated egg yolk samples. Total cholesterol, HDL cholesterol, total triglycerides concentrations of egg yolk were determined using RANDOX® cholesterol assay kit.

Statistical analysis

All data were subjected to analysis of variance using the General Linear Model Procedure SAS software (SAS, 2002). Treatment means were separated using the Duncan multiple range test at $p < 0.05$.

RESULTS

The effects of humic acid supplementation on the growth performance of ISA Brown and Fulani

ecotype chickens (FEC) are shown in Table 2. Compared with the control group, the supplementation of humic acid resulted in significant ($p < 0.05$) increase in body weight of ISA Brown and FEC at 8 and 20 weeks of age. Irrespective of the level of humic acid supplementation, ISA Brown chicks had similar body weight gain while, the body weight gain of FEC was significantly ($p < 0.05$) higher at 2 ml.L⁻¹ supplementation level.

Feed intake ($p < 0.05$) increased with humic acid supplementation in ISA Brown chicks, while it decreased in FEC at 8 weeks. However at 20 weeks of age, feed intake ($p < 0.05$) decreased in both ISA Brown and FEC with humic acid supplementation. Fulani ecotype chickens consumed ($p < 0.05$) less feed with or without humic acid supplementation when compared with the ISA Brown groups at 8 and 20 weeks.

Feed conversion ratio (FCR) was similar in all ISA Brown groups and FEC control group while the ratio decreased as the level of humic acid supplementation increased in FEC at 8 weeks. The highest FCR at 20 weeks was obtained from ISA Brown pullets without humic acid supplementation while FEC supplemented with 2 ml.L⁻¹ humic acid had the least value. Similar FCR values were recorded from the other treatment groups. Significant ($p < 0.05$) percentage mortality was recorded from ISA Brown and FEC without supplementation

Table 2: Growth performance of ISA Brown and Fulani ecotype chickens supplemented with humic acid

Parameter	ISA Brown			Fulani ecotype			SEM	p-value		
	Humic acid supplementation (mL.L ⁻¹)							HA	B	HAxB
	0	1	2	0	1	2				
Initial weight (g/bird)	33.17	33.19	33.16	33.19	33.19	33.16	0.01	0.1470	0.4103	0.4410
Final weight at 8 wks (g/bird)	562.34 ^{cd}	575.76 ^{ab}	578.26 ^{ab}	553.86 ^d	569.61 ^{bc}	584.69 ^a	2.76	0.0001	0.4026	0.0005
Weight gain at 8 weeks (g/bird)	529.18 ^{cd}	542.57 ^{ab}	545.10 ^{ab}	520.67 ^d	536.42 ^{bc}	551.53 ^a	2.76	0.0001	0.4011	0.0005
Feed intake at 8 wks (g/bird)	1729.26 ^b	1761.52 ^a	1764.53 ^a	1685.04 ^c	1648.00 ^d	1629.80 ^d	13.01	0.7723	0.0001	0.4011
Feed conversion ratio	3.27 ^a	3.25 ^a	3.24 ^a	3.24 ^a	3.07 ^b	2.96 ^c	0.03	0.0067	0.0003	<.0001
Mortality at 8 wks (%)	6.67 ^a	0.00 ^b	0.00 ^b	3.33 ^{ab}	0.00 ^b	0.00 ^b	0.90	0.0252	0.4752	0.1357
Body weight at 20 wks (g/bird)	1383 ^c	1500 ^a	1500 ^a	1410 ^{bc}	1463 ^{ab}	1490 ^a	13.73	0.0018	0.7344	0.0164
Weight gain at 20 wks (g/bird)	820.99 ^b	924.24 ^a	921.74 ^a	856.14 ^{ab}	893.73 ^{ab}	905.30 ^a	12.15	0.0118	0.8429	0.0589
Feed intake 8-20 wks (g/bird)	6399.41 ^a	6270.48 ^b	6251.14 ^b	5869.96 ^c	5603.00 ^d	5558.20 ^d	81.16	<.0001	<.0001	<.0001
Feed conversion ratio	7.82 ^a	6.79 ^b	6.78 ^b	6.86 ^b	6.27 ^{bc}	6.16 ^c	0.15	0.0005	0.0004	0.0006

^{a,b,c} Means on the same row with different superscript are different ($p < 0.05$)

SEM = Standard error of mean

HA = Humic acid, B = Breed

alone; there was no mortality among ISA Brown and FEC supplemented with humic acid.

The pH of the gastrointestinal tract (GIT) of ISA Brown and FEC supplemented with humic acid as indicated in Table 3 showed that the GIT was significantly ($p < 0.05$) affected by humic acid supplementation and breed. The pH of the crop, gizzard, duodenum, jejunum and ileum of ISA Brown and FEC ($p < 0.05$) reduced with humic acid supplementation. The ($p < 0.05$) highest pH values in each of the GIT segment were recorded in ISA Brown chicks in the control group while, FEC supplemented with 2 ml.L⁻¹ had the ($p < 0.05$) least pH values in these GIT segments.

Age at first lay, hen-day production, feed intake and egg weight were significantly ($p < 0.05$) improved

with humic acid supplementation in ISA Brown and FEC as revealed in Table 4. Highest ($p < 0.05$) hen-day production was obtained with FEC supplemented with humic acid while, the least was recorded from ISA Brown chickens without supplementation. Similar hen-day production percentages were obtained from the other treatment groups. Yolk percentage was significantly ($p < 0.05$) higher and similar in all FEC groups and ISA Brown not supplemented with humic acid. This was ($p < 0.05$) lower in ISA Brown chickens supplemented with humic acid.

Shell thickness was significantly ($p < 0.05$) increased at 2 ml.L⁻¹ humic acid supplementation level in ISA Brown and FEC although, shell thickness from ISA Brown chicken groups was ($p < 0.05$) higher than

Table 3: Gastrointestinal pH of ISA Brown and Fulani ecotype chickens supplemented with humic acid

Parameter	ISA Brown			Fulani ecotype			p-value			
	Humic acid supplementation (mL.L ⁻¹)						SEM	Humic	Breed	HAXB
	0	1	2	0	1	2				
Crop	5.17 ^a	5.01 ^b	4.87 ^c	4.41 ^d	4.23 ^e	4.15 ^e	0.10	<.0001	<.0001	<.0001
Gizzard	4.87 ^a	4.75 ^b	4.68 ^c	4.16 ^d	3.66 ^e	3.38 ^f	0.14	0.0352	<.0001	<.0001
Duodenum	5.71 ^a	5.58 ^b	5.47 ^c	5.45 ^{cd}	5.42 ^d	5.42 ^d	0.03	0.0020	<.0001	<.0001
Jejunum	5.84 ^a	5.72 ^b	5.63 ^c	5.59 ^c	5.46 ^d	5.41 ^d	0.04	<.0001	<.0001	<.0001
Ileum	6.55 ^a	6.32 ^b	6.22 ^b	6.03 ^c	5.92 ^{cd}	5.83 ^d	0.06	0.0001	<.0001	<.0001

^{a,b,c} Means on the same row with different superscript are different ($p < 0.05$)

SEM = Standard error of mean

HA = Humic acid, B = Breed

Table 4: Production performance and egg qualities of ISA Brown and Fulani ecotype chickens supplemented with humic acid

Parameter	ISA Brown			Fulani ecotype			p-value			
	Humic acid supplementation (mL.L ⁻¹)						SEM	Humic	Breed	HAXB
	0	1	2	0	1	2				
Age at 1 st egg (days)	153 ^a	146 ^b	145 ^b	142 ^c	139 ^d	139 ^d	1.17	0.0003	<.0001	<.0001
Hen-day production (%)	87.33 ^c	92.67 ^b	92.00 ^b	92.67 ^b	95.33 ^a	95.33 ^a	0.68	<.0001	<.0001	<.0001
Feed intake (g/bird/day)	113.00 ^a	110.00 ^b	109.00 ^b	91.67 ^c	90.00 ^{cd}	89.67 ^d	2.48	0.0007	<.0001	<.0001
Egg weight (g/egg)	56.37 ^c	59.34 ^b	64.75 ^a	40.52 ^f	43.56 ^c	46.17 ^d	2.16	<.0001	<.0001	<.0001
Yolk (%)	29.06 ^a	27.49 ^b	27.02 ^b	28.90 ^a	29.19 ^a	29.59 ^a	0.25	0.2832	0.0025	0.0002
Shell (%)	9.49 ^b	9.41 ^b	9.30 ^b	9.86 ^a	9.57 ^{ab}	9.44 ^b	0.06	0.0443	0.0244	0.0570
Shell thickness (mm)	0.33 ^{bc}	0.34 ^b	0.35 ^a	0.32 ^d	0.32 ^d	0.33 ^c	0.03	0.0021	<.0001	0.0002
Egg shape index	0.67 ^b	0.67 ^b	0.73 ^a	0.67 ^b	0.65 ^c	0.74 ^a	0.01	<.0001	0.3262	<.0001
Haugh Unit (%)	83.06 ^b	84.12 ^a	84.30 ^a	78.75 ^c	79.26 ^c	78.92 ^c	0.61	0.0720	<.0001	<.0001
Yolk colour	8.00 ^c	9.00 ^b	10.00 ^a	6.00 ^e	7.00 ^d	7.00 ^d	0.33	<.0001	<.0001	<.0001

^{a,b,c} Means on the same row with different superscript are different ($p < 0.05$)

SEM = Standard error of mean

HA = Humic acid, B = Breed

those from the FEC groups. Haugh unit ($p < 0.05$) improved with humic acid supplementation in ISA Brown chickens while, there was no significant ($p > 0.05$) effect of humic acid on all FEC groups. Haugh unit was significantly ($p < 0.05$) higher in ISA Brown chickens when compared with FEC. Yolk colour was significantly ($p < 0.05$) higher in the groups supplemented with humic acid when compared to those in the control groups. ISA Brown chickens supplemented with 2 mL.L⁻¹ produced eggs with deeper yolk colour, which was closely followed by the ISA Brown group supplemented with 1 mL.L⁻¹ humic acid. The ($p < 0.05$) lowest yolk colour was obtained from FEC in the control group.

Table 5 showed the egg lipid profile of ISA Brown and FEC supplemented with humic acid.

Triglycerides ($p < 0.05$) decreased in the eggs of ISA Brown and FEC supplemented with humic acid. However, egg cholesterol ($p < 0.05$) increased with humic acid supplementation. The highest cholesterol value was obtained from ISA Brown supplemented with 2 mL.L⁻¹ humic acid and the least from FEC supplemented with 2 mL.L⁻¹ humic acid. High density lipoprotein ($p < 0.05$) increased and decreased across the treatment groups in ISA Brown and FEC respectively. Low density lipoprotein ($p < 0.05$) increased with humic acid supplementation in ISA Brown and FEC. The highest low density lipoprotein value was recorded from ISA Brown chickens supplemented with 2 mL.L⁻¹ humic acid while, the least value was obtained from FEC without humic acid supplementation.

Table 5: Lipid profile of eggs from ISA Brown and Fulani ecotype chickens supplemented with humic acid

Parameter	ISA Brown			Fulani ecotype			p-value			
	Humic acid supplementation (mL.L ⁻¹)						SEM	Humic	Breed	HAXB
	0	1	2	0	1	2				
Triglycerides (mg.g ⁻¹)	151.74 ^a	117.99 ^b	112.39 ^{bc}	94.46 ^{bc}	87.70 ^c	87.90 ^c	6.11	0.0402	0.0001	0.0008
Cholesterol (mg.g ⁻¹)	75.36 ^d	77.50 ^c	79.28 ^a	74.64 ^d	78.32 ^b	70.98 ^e	0.68	0.0777	0.0240	<.0001
High density lipoprotein (mg.g ⁻¹)	44.13 ^c	45.59 ^b	47.67 ^a	39.66 ^d	30.57 ^e	32.52 ^f	1.57	0.1018	<.0001	<.0001
Low density lipoprotein (mg.g ⁻¹)	24.65 ^c	25.64 ^b	26.65 ^a	21.47 ^e	23.61 ^d	24.44 ^c	0.40	<.0001	<.0001	<.0001

^{a,b,c} Means on the same row with different superscript are different ($p < 0.05$)

SEM = Standard error of mean

HA = Humic acid, B = Breed

DISCUSSION

The improved body weight gain of birds supplemented with humic acid when compared with the un-supplemented group could be due to the beneficial effect of humic acid on the gut flora. The beneficial microbiological and pH-decreasing abilities of humic acid especially at the upper part of the gastrointestinal tract might have had resulted in the inhibition of intestinal bacteria leading to the reduced metabolic needs, thereby increasing the availability of nutrients to the host. Humic acid stimulates the immune system receptors in the gut lining to protect against pathogens to promote growth (Kocabagli *et al.*, 2002; Karaoglu *et al.*, 2004). By modifying the intestinal pH, Owing *et al.* (1990) reported that organic acid improves the solubility of the feed ingredients, digestion and absorption of nutrients. Sheikh *et al.* (2010) revealed that organic acid supplementation facilitated nutrient absorption to a greater extent and thus boosted growth.

The acid anion of humic acid has been shown to complex with calcium, phosphorus, magnesium and zinc which results in an improved digestibility of these minerals and serves as substrate in the intermediary metabolism (Kishi *et al.*, 1999).

The increase in feed intake of ISA Brown chicks and decrease in FEC with humic acid supplementation may be as a result of the differences in breed. Kucukersan *et al.* (2005) showed that the average daily feed consumption of hen fed diets with humic acid was significantly decreased compared with the control group. While some researchers reported that dietary humate supplementation increased feed intake (Hayirli *et al.*, 2005). The improvement in the FCR among FEC supplemented with humic acid could be possibly due to better utilization of nutrients resulting in increased body weight. Supplementation of humic acid at levels of 1 and 2 g.kg⁻¹ were reported to improve feed efficiency when compared with hens fed control diet (Yoruk *et al.*, 2004). Organic acids improve the absorption and conversion of nutrients in the body, and improve overall gastric function (Park *et al.*, 2009).

In this study, humic acid was able to improve the survivability of the birds; this could be attributed to the reduction in the intestinal pH and the ability to inhibit the growth of microbial pathogens. Humic acid was observed to show antibacterial, antiviral, antithyroidal and anti-inflammatory effects in animals, thus improves the immune system (Islam *et al.* 2005).

Humic acid has the ability to influence in particular the metabolism of protein carbohydrates of microbes by catalytic means. This leads to a direct devastation of bacterial cells or virus particles (Huck *et al.*, 1991; Ricke, 2003).

The use of humic acids in animal feeds improved

animal health forming a protective film on the mucous epithelium of the membrane and tract against infection and toxin (Kucukersan *et al.*, 2005). The lower pH values in GIT of birds supplemented with humic acid is a confirmation of the pH decreasing ability of humic acid. The addition of acidifiers to the diet for broilers lowered the pH of the crop and gizzard content (Andrys *et al.*, 2003).

Egg production and egg weight among the groups supplemented with humic acid is indicative of better utilization of nutrients. Egg production values of laying hens fed 1 and 2 g.kg⁻¹ humic acid were higher than those of the control group (Tanco, 1999; Yoruk *et al.*, 2004). However, some studies found that humic acid had no effect on egg production of laying hens and laying quails (Kucukersan *et al.*, 2005; Yalcin *et al.*, 2005). Soltan (2008) found that organic acid supplementation increased egg production by about 5.77 % compared to untreated group. Park *et al.* (2002) reported a positive effect of organic acid on laying performance. Dietary humic substances improved egg weight as reported by Wang *et al.* (2007).

The increased egg shell quality in eggs produced by birds supplemented with humic acid might be as a result of the presence of calcium in humic acid and increase in calcium absorption. Increased permeability allowed easier transfer of minerals from the blood to the bone and cells (Enviromate, 2002). Organic acid is reported to improve utilization of mineral feed additives, increasing the availability of calcium and other minerals (Omogbenigun *et al.*, 2003). The pH reducing and antimicrobial effects of the organic acid may assist in gut acidifiers and appear to be a good solution for poor egg shell quality (Dibner and Butin, 2002; Dhawale, 2005). Thick egg shell is essential for protection against the penetration of pathogenic bacteria (Swiatkiewicz *et al.*, 2010).

Egg lipid profile showed certain interaction between breeds and their capability to take up dietary nutrients and transfer them to their products. Sarica *et al.* (2009) reported higher cholesterol in the eggs of commercial layers and lower in local pure breeds. However, Rizzi and Chiericato, (2010) revealed that yolk cholesterol was found at higher concentration in local Mediterranean laying hen breeds compared with commercial laying hybrids.

CONCLUSION

Supplementing ISA Brown and Fulani ecotype chickens with humic acid increased body weight, egg production, egg weight, shell thickness and improved feed conversion. Humic acid at the rate 1 mL.L⁻¹ can be used for production efficiency in ISA Brown and Fulani ecotype chickens.

ACKNOWLEDGEMENT

The authors are grateful to the Institute of Food Security, Agricultural Research and Environmental Resources for the provision of the Fulani ecotype chickens to carry out this study under the Fulani ecotype multiplication project.

REFERENCES

- ADEBAMBO, O. A. – IKEOBI, C. O. N. – OZOJE, M. O. – ADEBAMBO, O. A. 2009. Variation in growth performance of pure and crossbred meat type chickens. *Nigerian Journal of Animal Production*, vol. 36 (2), 2009, p. 211–227.
- ANDRYS, R. – KLECKER, D. – ZEMAN, L. – MAREČEK, E. 2003. The effect of changed pH values of feed in isophosphoric diets on chicken broiler performance. *Czech Journal of Animal Science*, vol. 48, 2003, p. 197–206.
- AOAC. 1990. *Official Methods of Analysis*. 14th edn. Association of Official Analytical Chemists. Arlington, Virginia.
- DHAWALE, A. 2005. Better eggshell quality with a gut acidifier. *Poultry International*, vol. 44, 2005, p. 18–21.
- DIBNER, J. J. – BUTIN, P. 2002. Use of organic acids as a model to study the impact of gut microflora on nutrition and metabolism. *Journal of Applied Poultry Research*, vol. 11, 2002, p. 453–463.
- ENVIROMATE, T. M. 2002. Effects of humic acid on animals and humans (literature review and current research). Effects of humic acid. Enviromate Inc. 8571 Boat Club Road, Forth Worth, Texas 76179. <http://www.enviromateinc.com/effectsha.asp>
- EREN, M. – DENİZ, G. – GEZEN, S. S. – TURKMEN, I. I. 2000. Effects of humates supplemented to the broiler feeds on fattening performance, serum mineral concentration and bone ash. *Veterinary Journal of Ankara University*, vol. 47, 2000, p. 255–263.
- FAO, 2000. Food and Agricultural Organization small scale poultry production, FAO Corporate Document Repository, <http://www.fao.org/docrep/008/y5169e01.htm#>
- FAYEYE, T. R. – ADESHIYAN, A. B – OLUGBAMI, A. A. 2005. Egg Traits, Hatchability and Early Growth Performance of the Fulani – Ecotype Chicken. *Livestock Research for Rural Development*, vol. 17 (8), 2005, Art. #94. <http://www.lrrd.org/lrrd17/8faye17094.htm>.
- FMA&RD. 2006. Federal Ministry of Agriculture and Rural Development, Federal Department of Livestock and Pest Control Service. 'Nigeria Poultry Population' Highly pathogenic avian influenza standard operating procedure. February, 2006.
- GALIK, B. – ROLINEC, M. 2011. The effect of phytogenic feed additives on the performance of non-ruminant Proc. International Ph.D. workshop on welfare, biotechnology and quality of animal production, Nitra, Slovak University of Agriculture, vol. 19, 2011.
- HAUGH, R. R. 1937. The Haugh unit for measuring egg quality. *US egg and poultry magazine*, vol. 43, 1937, p. 552–555.
- HAYIRLI, A. – ESENBUGA, N. – MACIT, M. – LACIN, E. – KARAOGLU, M. – KARACA, H. – YILDIZ, L. 2005. Nutrition practice to alleviate the adverse effects of stress on laying performance, metabolic profile, and egg quality in peak producing hens: I. The humate supplementation. *Asian-Australian Journal of Animal Science*, vol. 18, 2005, p. 1310–1319.
- HORST, P. 1989. Native fowl as reservoir for genome and major genes with direct and indirect effects on adaptability and their potential for tropical oriented breeding plans. *Archiv für Geflügelkunde*, vol. 53, 1989, p. 93–101.
- HUCK, J. A. – PORTER, N. – BUSHELD, M. E. 1991. Effect of humates on microbial activity. *Journal of General Microbiology*, vol. 137, p. 2321–2329.
- ISLAM, K. M. – SCHUHMACHER, A. – GROPP, J. M. 2005. Humic acid substances in animal agriculture. *Pakistan Journal of Nutrition*, vol. 4, 2005, p. 126–134.
- KARAOGLU, M. – MACIT, M. – ESENBUGA, N. – DURDAG, H. – TURGUT, L. – BILGIN, O. C. 2004. Effect of supplemental humate at different levels on the growth performance slaughter and carcass traits of broilers. *International Journal of Poultry Science*, vol. 3, 2004, p. 406–410.
- KIRCHGEßNER, M. – ROTH, F. X. 1982. Fumaric acid as a feed additive in pig nutrition. *Pig News Information*, vol. 3, 1982, p. 259–264.
- KISHI, M. – FUKAYA, M. – TSUKAMOTO, Y. – NAGASAWA, T. – KAKEHANA, K. – NISHIZAWA, N. 1999. Enhancing effect of dietary vinegar on the intestinal absorption of calcium in ovariectomized rats. *Bioscience Biotechnology Biochemistry*, vol. 63, 1999, p. 905–910.
- KOCABAGLI, N. – ALP, M. – ACAR, N. – KAHRAMAN, R. 2002. The effects of dietary humate supplementation on broiler growth and carcass yield. *Poultry Science*, vol. 81, 2002, p. 227–230.
- KOPECKY, J. – HRNCAR, C. – WEIS, J. 2012. Effect of Organic Acids Supplement on Performance of Broiler Chickens. *Animal Sciences and Biotechnologies*, vol. 45 (1), 2012, p. 51–54.
- KRAL, M. – ANGELOVICOVA, M. – MRAZOVA, I. – TKACOVA, J. – KLIMENT, M. 2011. Probiotic and acetic acid of broiler chickens performance. *Scientific Papers Animal Science and Biotechnologies*, vol. 44, 2011, p. 149–152.

- KUCUKERSAN, S. – KUCUKERSAN, K. – COLPAN, I. – GONCUOGLU, E. – REISLI, Z. – YESILBAG, D. 2005. The effects of humic acid on egg production and egg traits of laying hen. *Veterinarni Medicina*, vol. 50 (9), 2005, p. 406–410.
- OLAWUNMI, O. O. – SALAKO, A. E. – AFUWAPE, A. A. 2008. Morphometric and differentiation and assessment of function of the Fulani and Yoruba ecotype indigenous chickens of Nigeria. *International Journal of Morphology*, vol. 26, 2008, p. 975–980.
- OMOGBENIGUN, F. O. – NYACHOTI, C. M. – SLOMINSKI, B. A. 2003. The effect of supplementing microbial phytase and organic acids to a corn-soybean based diet fed to early weaned pigs. *Journal of Animal Science*, vol. 81, 2003, p. 1806–1813.
- OWINGS, W. J. – REYNOLDS, D. L. – HASIAK, R. J. – FERKET, P. R. 1990. Influence of dietary supplementation with *Streptococcus faecium* M-74 on broiler body weight feed conversion; carcass characteristics and intestinal microbial colonization. *Poultry Science*, vol. 69, 1990, p. 1257–1264.
- PARK, J. H. – PARK, G. G. – RYU, K. S. 2002. Effect of feeding organic acid mixture and yeast culture on performance and egg quality of laying hens. *Korean Journal of Poultry Science*, vol. 29, 2002, p. 109–115.
- PARK, K. W. – RHEE, A. R. – UM, J. S. – PAIK, I. K. 2009. Effect of dietary available phosphorus and organic acids on the performance and egg quality of laying hens. *Journal of Applied Poultry Research*, vol. 18, 2009, p. 598–604.
- PETRUSKA, P. – TUSIMOVA, E. – KALAFOVA, A. – HASCIK, P. – KACANIOVA, M. – KOLESAROVA, A. – CAPCAROVA, M. 2012. Effect of propolis in chicken diet on selected parameters of mineral profile. *Journal of Microbiology, Biotechnology and Food Sciences*, vol. 1, 2012, p. 1090–1097.
- RICKE, S. C. 2003. Perspectives on the use of organic acids and short chain fatty acids as antimicrobials. *Poultry Science*, vol. 82 (4), 2003, p. 632–639.
- RIZZI, C. – CHIERICATO, G. M. 2010. Chemical composition of meat and egg yolk of hybrid and Italian breed hens reared using an organic production system. *Poultry Science*, vol. 89, 2010, p. 1239–1251.
- SARICA, M. – SEKEROGU, A. – KARACAY, N. 2009. Effect of genotype on fatty acid and cholesterol contents of hen's egg. *Asian Journal of Chemistry*, vol. 21, 2009, p. 511–516.
- SAS. 2002. SAS/STAT Statistical Analytical Systems User's guide (Release 9.1 Edition). SAS Institute Inc. Cary, North Carolina.
- SHEIKH, A. – TUFAIL, B. – GULAM, A. B. – MASOOD, S. M. – MANZOOR, R. 2010. Effect of Dietary Supplementation of Organic Acids on Performance, Intestinal Histomorphology, and Serum Biochemistry of Broiler Chicken. *Veterinary Medicine International*, 2010, Article ID 479485, 7 pages <http://dx.doi.org/10.4061/2010/479485>
- SHEIKH, A. – TUFAIL, B. – GULAM, A. B. – MASOOD, S. M. – MASHUQ, R. – SYED, S. 2011. Response of broiler chicken to dietary supplementation of organic acids. *Journal of Central European Agriculture*, vol. 12 (3), 2011, p. 498–508.
- SOLTAN, M. A. 2008. Effect of organic acid supplementation on egg production, egg quality, and some blood serum parameters in laying hens. *International Journal of Poultry Science*, vol. 7, 2008, p. 613–621.
- SWIATKIEWICZ, S. – KORELESKI, J. – ARCZEWSKA, A. 2010. Laying performance and egg shell quality in laying hens fed diets supplemented with prebiotics and organic acids. *Czech Journal of Animal Science*, vol. 55 (7), 2010, p. 294–306.
- TANCHO, A. 1999. The effect of humic acid supplement on egg performance and egg quality of laying hen during 28-47 weeks of age. 30th Anniversary Kaset Ladkrabang, 24-26 June 1999. <http://www.kmitl.ac.th/agritech30/example.doc>
- WANG, Q. – KIM, H. J. – CHO, J. H. – CHEN, Y. J. – YOO, J. S. – KIM, I. H. 2007. Effects of supplemental humic substances on egg production and quality in laying hens. *Poultry Science*, vol. 86, p. 519.
- YALCIN, S. – ERGUN, A. – EROL, H. – YALCIN, S. – OZSOY, B. 2005. Use of L-carnitine and humate in laying quail diets. *Acta Veterinaria Hungarica*, vol. 53, 2005, p. 361–370.
- YILDIZ, G. – SACAKLI, P. – GUNGOR, T. 2006. The effect of dietary Jerusalem artichoke (*Helianthus tuberosus*) on performance, egg quality characteristics and egg cholesterol content in laying hens. *Czech Journal of Animal Science*, vol. 8, 2006, p. 349–354.
- YORUK, M. A. – GUL, M. – HAYIRLI, A. – MACIT, M. 2004. The effects of supplementation of humate and probiotic on egg production and quality parameters during the late laying period in hens. *Poultry Science*, vol. 83, 2004, p. 84–88.

EFFECTS OF SEASONAL CHANGES ON THE NUTRITIVE QUALITY OF *MORINGA OLEIFERA* AND *PANICUM MAXIMUM* SILAGE

V. O. A. OJO^{1*}, E. D. OYEBANJO¹, O. T. OPAYEMI¹, S. O. MUSTAPHA¹, O. O. ADELUSI², S. A. ADEOYE¹, U. Y. ANELE³, A. O. OGUNSAKIN¹, A. O. JOLAOSHO¹, O. S. ONIFADE¹

¹Department of Pasture and Range Management, Federal University of Agriculture, Abeokuta, Ogun State, Nigeria

²Department of Animal Nutrition, Federal University of Agriculture, Abeokuta, Ogun State, Nigeria

³Agriculture and Agri-Food Canada Lethbridge Research Centre, Lethbridge, Alberta, Canada

ABSTRACT

The experiment was carried out to evaluate the effects of season and plant types on the chemical composition and *in vitro* fermentation of *Moringa oleifera* and *Panicum maximum* (local) silage. Leaf samples of *M. oleifera* plants and *P. maximum* were randomly collected during the rainy and dry seasons from Federal University of Agriculture, Abeokuta, Nigeria. The design of this study was 2 × 5 factorial arrangements with 3 replicates (2 seasons and 5 mixture ratios). The forage samples were both chopped into 2–3 cm lengths and divided into different proportions of 100 % : 0 %, 0 % : 100 %, 30 % : 70 %, 50 % : 50 % and 70 % : 30 % and were carefully packed into 30 laboratory bottles (960 ml) silos after mixing thoroughly. Season × plant species interactions were observed for some chemical and *in vitro* fermentation parameters. The CP content of the samples was significantly higher ($p < 0.05$) and ranged from 69.9 to 165.5 g/kg DM⁻¹ and fibre concentrations ranged from 246.3 to 710.5 g/kg DM⁻¹, 193.2 to 578.0 g/kg DM⁻¹ and 63.5 to 133.3 g/kg DM⁻¹ for NDF, ADF and ADL respectively. 100 % Moringa + 0 % Panicum silage recorded the highest ($p < 0.05$; 165.5 g/kg DM⁻¹) CP content in all the silages evaluated. All the silages had high calcium contents (7.0–11.7 g/kg DM⁻¹). Values obtained for the *in vitro* fermentation characteristics of the forage species indicated the presence of potentially degradable nutrients. 100 % Moringa + 0 % Panicum and all the forage mixtures silages have potentials that could be harnessed as feed supplements for ruminants both in the rainy and dry seasons.

Key words: Moringa; nutritive value; Panicum; plant types; season; silage

INTRODUCTION

Forages provide adequate amount of nutritious feed for ruminants in the form of grasses which is preferred by these types of animals. However, in the dry season, there is feed constraint due to seasonal shortages in the quantity and quality of forage from natural pastures that provide most of the feed for animals which result into problems such as sickness and weight loss due to poor dietary profile for the animals and reduction in yield and quality of forages arises. Under this condition, browse plant is of great importance in balancing protein deficiency during the long dry season. They are effective insurance against seasonal feed shortages. The main

feature of browse plants is their high crude protein (CP) and mineral contents (Awad and Elhadi, 2010).

Ruminants relish guinea grass (Babayemi and Bamikole, 2006). But such forage becomes very scarce in the dry season. It is commonly used for silage but herbage has been reported to have low content of dry matter and low concentrations of water soluble carbohydrates and crude protein (Zanine *et al.*, 2006). The levels of protein content and fermentable carbohydrates can be improved through various treatments including mixing legumes with cereal crops (Phiri *et al.*, 2007).

Browse plants have special characteristics of high nitrogen value, better metabolizable energy

***Correspondence:** E-mail: ojovoa@funaab.edu.ng
 Victoria Olubunmi A. Ojo, Department of Pasture and Range Management,
 Federal University of Agriculture, P. M. B. 2240, Abeokuta, Ogun State, Nigeria
 Tel.: +2348037178044

Received: May 5, 2015

Accepted: June 2, 2016

and organic matter digestibility when fed with grass supplement (Ajayi and Babayemi, 2008). Forage tree leaves generally have high phosphorus concentrations (McMeniman and Little, 1974).

In dealing with lack of feed during the dry season forage conservation especially as silage is considered to be the preservation technique with the greatest potential for protein rich foliage (Man and Wiktorsson, 2002). Ensiling cereal and legumes together have been reported to be responsible in providing sufficient fermentable carbohydrates for lactic acid bacteria (Koljajic *et al.*, 1998) and simultaneously the protein content of silage will be increased (Anele *et al.*, 2011). In addition, mixing legumes and grasses increases biomass yield, crude protein (CP) content, nutritive value of resultant silage and soil fertility (Assefa and Ledin, 2001; Nayigihugu *et al.*, 2002).

This study will therefore evaluate the chemical composition and *in vitro* fermentation of *Moringa oleifera* and *Panicum maximum* silage at different proportions and seasons.

MATERIAL AND METHODS

Experimental site

The experiment was conducted at the Department of Pasture and Range Management, University of Agriculture, Abeokuta, Nigeria. The site lies within the derived savannah zone of South Western Nigeria on latitude 7°58'N, longitude 3°20'E and 75 m above sea level with average annual rainfall of 1037 mm. Abeokuta has a bimodal rainfall pattern that typically peaks in July and September with a break of two to three weeks in August. Temperatures are fairly uniform with daytime values of 28 to 30 °C during the rainy season (April – September) and 30–34 °C during the dry season (October – March) with the lowest night temperature of around 24 °C during the harmattan period between December and February. Relative humidity is high during the rainy season with values between 63 and 96 % as compared to dry season 55–84 %. The temperature of the soil ranges from 24.5 to 31.0 °C (Source: Agromet Dept., FUNAAB).

Sample collection

P. maximum samples were harvested at 15 cm above ground level at 6 weeks after cut back from the Fadama (wetland areas) located within the University and *M. oleifera* was also harvested by hand cutting the branches followed by stripping off all leaves in July, 2011 for rainy season and December, 2011 for dry season from the Teaching and Research Farm of the University. The grass samples were chopped into pieces of 2-3 cm in length, wilted for 4 hours to reduce their moisture contents

at a temperature of 23.4 °C and relative humidity of 71.2 % before ensiling and divided into different proportions of 100 %:0 %, 0 %:100 %, 30 %:70 %, 50 %:50 % and 70 %:30 %.

Experimental design

The experiment was laid out in a 2 x 5 factorial design arrangement with 3 replicates consisting of two types of season i.e. rainy and dry, and five forage proportions i.e. 100 % Moringa + 0 % Panicum, 0 % Moringa + 100 % Panicum, 50 % Moringa: 50 % Panicum, 70 % Moringa: 30 % Panicum, and 30 % Moringa:70 % Panicum.

Data collection

The chopped and wilted plant types at different forage proportions and their sole were carefully packed into laboratory (960 ml) bottle silos after mixing thoroughly following the method described by Yokota *et al.* (1995). A total of thirty (30) bottle silos were used in the ensiling process. The forage samples were ensiled for a period of 6 weeks at an ambient temperature of 26 °C. At the expiration of the ensiling, the bottle silos were opened, the pH of the silages was determined and chemical and *in vitro* analyses of the samples were thereafter carried out.

Chemical analyses

Samples of 300 g were taken from each of the bottle silos and oven-dried to a constant weight at 65 °C. The dried foliage samples were milled through a 1mm sieve and crude protein (CP), ether extract (EE), ash and the minerals such as Phosphorus (P), Potassium (K), Calcium (Ca) and Sodium (Na) were analyzed according to the standard methods of AOAC (2000). Neutral detergent fibre (NDF), acid detergent fibre (ADF) and acid detergent lignin were determined according to Van Soest *et al.* (1991). Hemicellulose was calculated as NDF – ADF while Cellulose was calculated as ADF – ADL. Non-fibre carbohydrates (NFC) were calculated as:

$$\text{NFC} = 1000 - \text{CP} - \text{ash} - \text{EE} - \text{NDF}$$
with all variables expressed as g/kg DM⁻¹.

In vitro gas production was determined according to Menke and Steingass (1988). Rumen fluid was collected from three bulls before morning feeding according to the method described by Babayemi and Bamikole (2006). The animals have been previously fed a mixed diet of *P. maximum* grass and concentrates to fulfil maintenance requirements. The liquor was collected into a pre warmed thermos flasks and was later filtered through three layers of cheesecloth. Two hundred milligrams of ground feed samples were weighed into 100 ml calibrated syringes with pistons lubricated with grease. A buffered mineral solution

was prepared. 30 ml of buffered rumen fluid was taken into syringes containing the feeds. The syringes were positioned vertically in a water bath with shaker kept at 39 °C fitted with plungers. Gas production rates were recorded at 3, 6, 9, 12, 24, 36 and 48 hour of incubation and each syringe was gently swirled after reading. Three blank syringes containing only 30 ml of buffered inoculum were also concurrently incubated.

The data obtained were fitted to the non-linear regression equation:

$$V \text{ (ml.200 mg.DM}^{-1}\text{)} = b (1 - e^{-ct})$$

where V = potential gas production at time t , b = volume of gas that will evolve with time and c = fractional rate of gas production. Initial gas production rate (A_{bsg}) was calculated as the product of b and c (Larbi *et al.*, 1996).

Organic matter digestibility (OMD) were estimated as:

$$\text{OMD} = 14.88 + 0.889\text{GV} + 0.45 \text{ CP} + 0.651 \text{ ash (Menke and Steingass, 1988).}$$

Short-chain fatty acids (SCFA) were estimated as:

$$\text{SCFA} = 0.0239\text{GV} - 0.0601 \text{ (Getachew et al., 2000).}$$

Metabolizable energy (ME) were calculated as:

$$\text{ME} = 2.20 + 0.1357\text{GV} + 0.0057 \text{ CP} + 0.0002859 \text{ EE}^2 \text{ (Menke and Steingass, 1988).}$$

Total gas volume (GV) were expressed as ml/200 mg DM, CP and ash as g.kg DM⁻¹, ME as MJ.kg DM⁻¹ and SCFA as μmol.g DM⁻¹.

In vitro dry matter digestibility (IVDMD)

After 48 hours digestion, the bottle contents were centrifuged at 15,000 rpm for 10 minutes and residues were decanted into pre weighed crucibles were oven-dried at 105 °C for 24 hours. The dry residues were weighed and digestibility calculated using the equation as follows:

$$\text{IVDMD (\%)} = \frac{\text{Initial DM input} - \text{DM residue-Blank} \times 100}{\text{Initial DM input}}$$

Data were subjected to two-way analysis of variance using the SAS (2002) package while significant means were separated using Duncan's multiple range test ($p < 0.05$).

RESULTS

Chemical composition

The proximate composition and pH of the silages are shown in Table 1. There were significant differences ($p < 0.05$) in the DM, CP, EE, ash and NFC composition as a result of interactions between plant types and season. 100 % Moringa + 0 % Panicum silage had the lowest DM contents for both the rainy and dry seasons compared with others. Meanwhile, 100 % Moringa + 0 % Panicum silage during rainy season recorded the highest CP value (165.5 g.kg DM⁻¹) which was closely trailed behind by mixtures of 30 % Moringa : 70 % Panicum silage at rainy (129.4 g.kg DM⁻¹). 0 % Moringa + 100 % Panicum silage at dry season recorded the least value of EE (10.1 g.kg DM⁻¹). 100 % Moringa + 0 % Panicum and

Table 1: Effect of season and plant type on the composition and pH of the silage (g.kg DM⁻¹)

Season	Forage mixture	DM	CP	EE	ASH	NFC	pH
				(g.kg DM ⁻¹)			
Rainy	100 % Moringa + 0 % Panicum	910.00 ^b	165.50 ^a	32.00 ^{abc}	80.4 ^c	327.20 ^b	4.48
	0 % Moringa + 100 % Panicum	945.00 ^a	74.10 ^c	25.90 ^{bcd}	88.9 ^{abc}	100.60 ^e	4.64
	Moringa 50 %: Panicum 50 %	935.00 ^a	127.90 ^b	47.10 ^a	86.6 ^{abc}	172.30 ^{cd}	4.38
	Moringa 30 %: Panicum 70 %	945.00 ^a	129.40 ^b	31.20 ^{abc}	103.4 ^{ab}	144.70 ^d	4.49
	Moringa 70 % : Panicum 30 %	960.00 ^a	83.90 ^c	31.00 ^{abc}	109.2 ^a	197.20 ^{cd}	4.60
Dry	100 % Moringa + 0 % Panicum	910.00 ^b	127.00 ^b	37.60 ^{ab}	90.1 ^{abc}	499.30 ^a	4.20
	0 % Moringa + 100 % Panicum	950.00 ^a	69.90 ^c	10.10 ^d	108.10 ^{ab}	60.90 ^f	4.50
	Moringa 50 % : Panicum 50 %	940.00 ^a	82.00 ^c	20.90 ^{bcd}	93.10 ^{abc}	298.70 ^c	4.70
	Moringa 30 % : Panicum 70 %	935.00 ^a	89.90 ^c	15.40 ^{cd}	107.60 ^{ab}	241.60 ^c	4.63
	Moringa 70 % : Panicum 30 %	945.00 ^a	77.8 ^c	36.60 ^{ab}	84.10 ^c	410.00 ^a	4.80
	SEM	3.50	6.40	2.50	2.70	2.60	0.06

^{abcd}Means along the same column with different subscripts are significantly different ($p < 0.05$)

DM: Dry matter, CP: Crude protein, EE: Ether extract, NFC: Non Fibre Content

SEM: Standard error of mean

mixtures of 70 % Moringa: 30 % Panicum silages both during dry season recorded the highest ($p < 0.05$) overall NFC.

Table 2 shows the effects of plant types and season on the fibre composition of forages. 0 % Moringa + 100 % Panicum silage at rainy and dry seasons had the highest ($p < 0.05$) NDF while 100 % Moringa + 0 % Panicum silage at dry season had the least ($p < 0.05$) NDF, ADF and celluloses levels, respectively.

Table 3 shows the mineral composition of the silages. There were significant ($p < 0.05$) interactions between plant types, forage proportions and season for the phosphorus, potassium, calcium and sodium contents of the silages. Among the interaction of the mineral contents of the forages, P and Ca contents ranged from 2.2 to 3.6 g/kg DM⁻¹ and 7.0 to 11.70 g/kg DM⁻¹ respectively. Silages produced from legumes and different plant types had higher amounts of these minerals than those from grasses alone. Potassium contents of the forage mixtures during the rainy season were significantly ($p < 0.05$) higher than in the dry season.

In vitro digestibility

There were significant effects of plant types and season on the volume of gas produced at different times for the silages (Table 4). 100 % Moringa + 0 % Panicum and 70 % Moringa : 30 % Panicum silages both at dry season, recorded the highest ($p < 0.05$) gas production at 3 hr of incubation while 0 % Moringa + 100% Panicum and 70 % Moringa : 30 % Panicum silages during rainy season and 0 % Moringa + 100 % Panicum

silage at dry season respectively produced no gas at 3 hr of incubation. However, at 48 hr of incubation, 30% Moringa : 70 % Panicum silage at dry season had the highest gas production with 35 % while 50 % Moringa : 50 % Panicum silage at rainy season produced the least gas of 23 % at 48 hr.

Irrespective of plant types and season of ensiling, there was no significant difference ($p > 0.05$) on insoluble but degradable fraction (b) and fractional rate of gas production (c) of the ensiled forages (Table 5). However, 70 % Moringa : 30 % Panicum silage at dry season had the highest SCFA and ME values above all other plant types and season. SCFA content of the silages of different plant types was higher during dry season except for 0 % Moringa + 100 % Panicum silage. Similar trend was noticed with ME.

DISCUSSION

Chemical composition

Higher DM contents as obtained in this study have been reported to concentrate the water soluble carbohydrates and improve the effectiveness of lactate-producing bacteria, and thus improve the quality of silage (Anele *et al.*, 2011).

The CP contents of all the silages of plant types were all above the critical CP level of 8 % acceptable for ruminant performance (Norton, 1994) except the 0 % Moringa + 100 % Panicum silage which was similar to 7.8 % reported by Aganga and Tshwenyane, (2004) and lower

Table 2: Effect of season and plant type on the fibre composition of the silage (g/kg DM⁻¹)

Season	Forage mixture	NDF	ADF	ADL (g/kg DM ⁻¹)	HEM	CELL
Rainy	100 % Moringa + 0 % Panicum	394.90 ^d	347.70 ^{de}	91.70 ^b	47.20 ^d	256.00 ^{de}
	0 % Moringa + 100 % Panicum	710.50 ^a	420.20 ^{bcd}	133.30 ^a	290.30 ^a	286.90 ^d
	Moringa 50 %: Panicum 50 %	566.10 ^{bc}	467.50 ^b	63.50 ^{cd}	98.70 ^{cd}	404.00 ^{ab}
	Moringa 30 %: Panicum 70 %	591.30 ^b	463.70 ^b	66.20 ^{bcd}	127.60 ^{bc}	397.50 ^{abc}
	Moringa 70 %: Panicum 30 %	578.70 ^b	347.50 ^{de}	50.00 ^d	231.20 ^a	297.50 ^{cd}
Dry	100 % Moringa + 0 % Panicum	246.30 ^c	193.20 ^f	80.10 ^{bc}	53.00 ^d	119.90 ^f
	0 % Moringa + 100 % Panicum	751.40 ^a	578.00 ^a	130.70 ^a	173.50 ^b	447.30 ^a
	Moringa 50 %: Panicum 50 %	505.30 ^c	427.30 ^{bc}	121.30 ^a	78.10 ^{cd}	332.60 ^{bc}
	Moringa 30 %: Panicum 70 %	545.50 ^{bc}	384.10 ^{cde}	82.00 ^{bc}	165.10 ^b	302.00 ^{bc}
	Moringa 70 %: Panicum 30 %	391.50 ^d	317.00 ^e	133.30 ^a	74.50 ^{cd}	183.70 ^{ef}
	SEM	27.3	19.2	6.0	17.0	19.6

^{abcdef} Means along the same column with different subscripts are significantly different ($p < 0.05$)

NDF: Neutral detergent fibre, ADF: Acid detergent fibre, ADL: Lignin detergent fibre, HEM: Hemicellulose

CELL: Cellulose

SEM = Standard error of mean

Table 3: Effect of season and plant type on the mineral content of the silage (g.kg DM⁻¹)

Season	Forage mixture	Phosphorus (P)	Potassium (K)	Calcium (Ca) (g.kg DM ⁻¹)	Sodium (Na)
Rainy	100 % Moringa + 0 % Panicum	3.56 ^a	33.26 ^c	11.72 ^c	2.90 ^e
	0 % Moringa + 100 % Panicum	2.44 ^g	30.57 ^e	7.02	2.71 ^g
	Moringa 50 %: Panicum 50 %	2.51 ^f	34.38 ^b	10.94 ^d	2.08 ⁱ
	Moringa 30 %: Panicum 70 %	2.23 ^h	31.89 ^a	8.52 ^f	2.53 ^h
	Moringa 70 % : Panicum 30 %	2.82 ^c	37.06 ^f	7.69 ^g	3.28 ^c
Dry	100 % Moringa + 0 % Panicum	3.05 ^b	31.39 ^d	10.48 ^a	3.13 ^d
	0 % Moringa + 100 % Panicum	2.51 ^f	24.26 ^h	7.76 ^g	2.83 ^f
	Moringa 50 % : Panicum 50 %	1.90 ⁱ	24.07 ^j	11.25 ^b	3.35 ^b
	Moringa 30 % : Panicum 70 %	3.05 ^b	29.71 ⁱ	11.70 ^c	4.52 ^a
	Moringa 70 % : Panicum 30 %	2.69 ^c	26.35 ^g	7.80 ^g	2.21 ⁱ
	SEM	0.08	1.39	0.71	0.12
Normal range* (%)		0.33-0.47	0.50-3.10	0.45-1.20	0.20-0.30

abcdehij Means along the same column with different subscripts are significantly different (p < 0.05)

SEM= Standard error of mean

* Normal range. Source: Rogers and Murphy (2000)

Table 4: Effect of season and plant type on the *in vitro* gas production of the silage (hr)

Season	Forage mixture	3	6	9	12	18	24	36	48
		(hr)							
Rainy	100 % Moringa	2.00 ^c	3.00 ^{dc}	8.00 ^c	10.00 ^d	12.00 ^d	17.00 ^e	23.00 ^f	25.00 ^f
	0 % Moringa + 100 % Panicum	0.00 ^e	2.00 ^{dc}	4.00 ^{dc}	7.00 ^e	10.00 ^e	19.00 ^d	28.00 ^{cd}	30.00 ^d
	Moringa 50 %: Panicum 50 %	1.00 ^d	4.00 ^{bc}	5.00 ^d	9.00 ^d	11.00 ^{dc}	15.00 ^f	20.00 ^g	23.00 ^g
	Moringa 30 %: Panicum 70 %	1.00 ^d	5.00 ^b	10.00 ^b	15.00 ^b	19.00 ^b	24.00 ^b	29.00 ^{cb}	33.00 ^{cb}
	Moringa 70 % : Panicum 30 %	0.00 ^e	3.00 ^{dc}	5.00 ^d	10.00 ^d	15.00 ^c	18.00 ^{dc}	28.00 ^{cd}	32.00 ^c
Dry	100 % Moringa + 0 % Panicum	4.00 ^a	7.00 ^a	10.00 ^b	15.00 ^b	19.00 ^b	21.00 ^c	26.00 ^c	28.00 ^c
	0 % Moringa + 100 % Panicum	0.00 ^e	1.00 ^c	3.00 ^c	7.00 ^c	12.00 ^d	18.00 ^{dc}	27.00 ^{cd}	34.00 ^{ab}
	Moringa 50 % : Panicum 50 %	3.00 ^b	5.00 ^b	10.00 ^b	15.00 ^b	19.00 ^b	24.00 ^b	30.00 ^{ab}	34.00 ^{ab}
	Moringa 30 % : Panicum 70 %	2.00 ^c	5.00 ^b	10.00 ^b	13.00 ^c	19.00 ^b	25.00 ^b	30.00 ^{ab}	35.00 ^a
	Moringa 70 % : Panicum 30 %	4.00 ^a	8.00 ^a	15.00 ^a	19.00 ^a	21.00 ^a	27.00 ^a	31.00 ^a	34.00 ^{ab}
	SEM	0.28	0.41	0.67	0.72	0.74	0.72	0.62	0.75

abcdehg Means along the same column with different subscripts are significantly different (P < 0.05)

than 9.17 % reported by Fadiyimu *et al.* (2000). Meanwhile, 100 % Moringa + 0 % Panicum silage CP contents for both seasons in this study were lower than 29.68 % reported by Fadiyimu *et al.* (2000). The differences between this study and the other authors could be because they used fresh forage sample in contrast to silage used in this study. Other factors could also be responsible for CP variation in samples. Wattiaux (1999) reported that proteolytic activities that takes place

during the ensilage could bring about reductions in CP content, while other variability could be attributed to plant parts, climatic condition, harvesting regime, location, soil type and age (Norton, 1994) and leaf to petiole ratio (Bamikole *et al.*, 2004). The lower CP content of 0 % Moringa + 100% Panicum silage compared to 100 % Moringa + 0 % Panicum silage in this study support the report of Mhere *et al.* (1999) that the major shortcoming of grasses is it low CP content. Hence the need for

Table 5: Effect of season and plant type on the fermentation kinetics of the silage

Season	Forage mixture	b (ml.200 mg DM ⁻¹)	c (ml.hr ⁻¹)	Lag time (hr)	SCFA (μ mol.g ⁻¹)	ME (MJ.kg ⁻¹)	OMD (%)	DMD (%)
Rainy	100 % Moringa + 0 % Panicum	33.14	0.03	1.60 ^{dc}	0.35 ^e	4.59 ^d	42.67 ^{cd}	70.00 ^{ab}
	0 % Moringa + 100 % Panicum	53.18	0.02	4.39 ^a	0.39 ^d	4.81 ^d	40.89 ^{ef}	50.00 ^e
	Moringa 50 %: Panicum 50 %	29.79	0.03	2.10 ^{dc}	0.30 ^f	4.30 ^e	39.61 ^f	60.00 ^{cd}
	Moringa 30 %: Panicum 70 %	36.86	0.05	2.53 ^{cb}	0.51 ^b	5.52 ^b	48.77 ^a	65.00 ^{cb}
	Moringa 70 % : Panicum 30 %	48.33	0.32	2.55 ^{cb}	0.37 ^{de}	4.68 ^d	41.76 ^{ef}	65.00 ^{cb}
Dry	100 % Moringa + 0 % Panicum	30.31	0.06	0.83 ^d	0.44 ^c	5.11 ^c	45.13 ^{cd}	65.00 ^{cb}
	0 % Moringa + 100 % Panicum	57.20	0.30	3.16 ^{ab}	0.37 ^{de}	4.67 ^d	41.06 ^{ef}	50.00 ^e
	Moringa 50 % : Panicum 50 %	39.20	0.04	1.65 ^{dc}	0.51 ^b	5.49 ^b	45.97 ^{cb}	55.00 ^{cd}
	Moringa 30 % : Panicum 70 %	41.09	0.04	2.19 ^{dc}	0.54 ^b	5.63 ^b	48.15 ^{ab}	75.00 ^a
	Moringa 70 % : Panicum 30 %	35.22	0.06	1.12 ^{dc}	0.59 ^a	5.89 ^a	47.86 ^{ab}	60.00 ^{cd}
	SEM	2.77	0.04	0.22	0.02	0.10	0.64	1.59

^{abcde}Means along the same column with different subscripts are significantly different ($P < 0.05$)

b: Insoluble but degradable fraction, c: Fractional rate of gas production, SCFA: Short chain fatty acid

ME: Metabolizable energy, OMD: Organic matter digestibility, DMD: Dry matter digestibility

mixture of grasses with browse plant that are high in CP, essential vitamin, minerals and amino acids is supported (Makkar and Becker, 1997, Gidamis *et al.*, 2003).

The range of non-fibre carbohydrates (NFC) content of the plant types silages in this study indicated that they can be easily degraded or fermented, as NFC is a crude estimate of the carbohydrate pool that differ in digestibility from NDF. The pH of silage is one of the simplest and quickest ways of evaluating its quality. The pH values of different ensiled forage mixtures at different season fell within 4.2 (Kung and Shaver, 2001) and 4.5 and 5.5 (Meneses *et al.*, 2007) that was recommended as the normal good silage for ruminants.

A higher CP content for 100 % Moringa + 0 % Panicum silage during the rainy season compared with dry season could be due to higher moisture content and nitrogen uptake being more rapid than dry matter accumulation during the rainy season. This agrees with Bamualim *et al.* (1980) and Larbi *et al.* (1997) who reported that seasonal variations occur between plant species and between seasons with higher crude protein values reported for seasons with higher moisture levels. The higher CP content of 100 % Moringa + 0 % Panicum silage in the rainy season could be due to the continuous flush (regrowth) of leaves during this period (Anele *et al.*, 2009).

0 % Moringa + 100 % Panicum silage recorded higher NDF, ADF and ADL above other plant types which was consistent with the observation of Okoli *et al.* (2003) that the fibre contents of tropical grasses is usually higher than that for browse, shrubs and trees. The range of NDF contents in this study for 100 %

Moringa + 0 % Panicum silage and all the forage mixtures for both seasons were below the range of 600-650 g.kg DM⁻¹ suggested as the limit above which intake of tropical feeds by ruminant animals would be limited (Van Soest *et al.*, 1991).

Silages produced from the legumes had higher Ca and P contents than those from the grasses. This is in accordance to the report of Nasrullah *et al.* (2004) and Marschner (1993) that there is a marked difference between the level of Ca and P in legumes and grasses. Since grasses have a tendency to be lower in Ca and P, hence the mixture of grasses and legumes in order to improve the nutrients available to animals. Ca is above the critical level of 3 g.kg DM⁻¹ as recommended for ruminants in the warm wet climates (McDowell *et al.*, 1993). This result indicates that Ca content of forage satisfied the animal requirements specified by the ARC (1980).

The P level in this study was above the critical level of 2.5 g.kg DM⁻¹ for ruminant animals as reported by Muhammad *et al.* (2005). The K contents in all plant type silages at different seasons of the year were higher than 8 g.kg DM⁻¹ as recorded for grazing animals (Underwood, 1981).

In vitro digestibility

The value of net gas produced in this study for 100 % Moringa + 0 % Panicum silage is similar to those reported for leaves from tropical trees and shrubs (*Adansonia digitata* and *Bombax glabra*) in south western Nigeria (Ogunbosoye and Babayemi, 2010). The low digestibility observed in this study for 100 %

Moringa + 0 % Panicum silage may be due to the presence of condensed tannins which is an anti-nutritional content found in most of the plant species (Barry and McNabb, 1999). Moringa (30 %) : Panicum (70 %) silage at dry season recorded the highest gas produced which suggests that it is of higher nutritional value than the other plant types and it connotes high digestibility of the forages at that mixtures.

The highest short-chain fatty acids (SCFA) and organic matter digestibility (OMD) contents in 70 % Moringa : 30 % Panicum silage in dry season and other forage mixtures, possibly because the mixtures contain more fermentable carbohydrate which is a vital substrate for growth of ruminal microorganisms (Van Soest, 1994). The estimation of the ME values is valuable for purposes of ration formulation and to set economic value of feed for other purposes (Getachew *et al.*, 2002). Gas production is a reflection of the generation of SCFA and microbial mass (Getachew *et al.*, 1998).

CONCLUSION

The pH values were within the range of pH for good quality silage. 100 % Moringa + 0 % Panicum silage recorded the highest CP content and lower fibre fractions than the 0 % Moringa + 100 % Panicum silage. Mixtures of the grass and browse plants produced quality silages for both the rainy and dry seasons. This will go a long way to provide quality feed for ruminants during the dry season when forages are scarce with low quality in the natural pastureland.

REFERENCES

- AGANGA, A. A. – TSHWENYANE, S. O. 2004. Potentials of guinea grass (*Panicum maximum*) as forage crop in livestock production. *Pakistan Journal of Nutrition*, vol. 3 (1), 2004, p. 1–4.
- AJAYI, F. T. – BABAYEMI, O. J. 2008. Comparative *in vitro* evaluation of mixtures of *Panicum maximum* cv Ntchisi with stylo (*Stylosanthes guianensis*), Lablab (*Lablab purpureus*), Centro (*Centrosema pubescens*) and hixrix (*Aeschynomene hixrix*). *Livestock Research for Rural Development*, vol. 20 (6), 2008, p. 1–15.
- ANELE, U. Y. – ARIGBEDE, O. M. – SÜDEKUM, K. -H. – ONI, A. O. – JOLAOSHO, A. O. – OLANITE, J. A. – ADEOSUN, A. I. – DELE, P. A. – IKE, K. A – AKINOLA, O. B. 2009. Seasonal chemical composition, *in vitro* fermentation and *in sacco* dry matter degradation of four indigenous multipurpose tree species in Nigeria. *Animal Feed Science and Technology*, vol. 154 (1), 2009, p. 47–57.
- ANELE, U. Y. – JOLAOSHO, A. O. – ARIGBEDE, O. M. – OLANITE, J. A. – ONIFADE O. S. 2011. Effects of air drying and plant types on quality of grass/legume silage. In: *Animal Feed: Types, Nutrition and Safety*. (Sarah R. Borgearo, ed). pp. 207–218. Nova Science Publishers, Inc. New York, USA.
- A.O.A.C. 2000: Association of Official Analytical Chemists. Official Methods of Analysis. 16th edition, Washington DC.
- ARC 1980. The nutrient requirements of ruminant livestock. pp. 309-314. Agricultural Research Council. (Commonwealth Agricultural Bureaux: Farnham Royal, Slough, G.B.).
- ASSEFA, G. – LEDIN, I. 2001. Effect of variety, soil type and fertiliser on the establishment, growth, forage yield, quality and voluntary intake by cattle of oats and vetches cultivated in pure stands and mixtures. *Animal Feed Science and Technology*, vol. 92, 2001, p. 95–111.
- AWAD, O. A – ELHADI, O. A. 2010: Seasonal variability in nutritive value of ruminant diets under open grazing system in the semi-arid rangeland of Sudan (South Darfur State). *Agriculture and Biology Journal of North America. ScienceHub*, vol. 1 (3), 2010, p. 243–249. <http://www.scihub.org/ABJNA>.
- BABAYEMI, O. J. – BAMIKOLE, M. A. 2006: Effects of Tephrosia candida DC leaf and its Mixtures with Guinea Grass on *in vitro* Fermentation Changes as Feed for Ruminants in Nigeria. *Pakistan Journal of Nutrition*, vol. 5, 2006, p. 14–18.
- BAMIKOLE, M. A. – IKHATUA, O. M. – ARIGBEDE, O. M. – BABAYEMI, O. J. – ETELA, I. 2004. An Evaluation of the Acceptability as forage of some Nutritive and Antinutritive Components and of the Dry Matter Degradation Profiles of Five Species of Ficus. *Tropical Animal Health and Production*, vol. 36, 2004, p. 157–167.
- BAMUALIM, A. – JONES, R. J. – MURRAY, R. M. 1980. Nutritive value of tropical browse legumes in the dry season. *Proceedings of the Australian Society of Animal Production*, 13, 1980, p. 229–232.
- BARRY, T. N. – MCNABB, W. C. 1999. The implication of condensed Tannins in the nutritive value of temperature forages fed to ruminants. *British Journal of Nutrition*, vol. 81, 1999, p. 263–272.
- FADIYIMU, A. A. 2000. Chemical composition, dry matter degradability and preference by WAD goats of some multipurpose trees in Nigeria. In: *Proceeding of 25th Annual Conference of Nigeria Society for Animal Production*, March 19–23, Umudike, Imo State, Nigeria. pp. 76–77.
- GETACHEW, G. – MAKKAR, H. P. S. – BECKER, K. 1998. The *in vitro* gas measuring techniques for assessment of nutritional quality of feeds: a review. *Animal Feed Science and Technology*,

- vol. 72, 1998, p. 261–281.
- GETACHEW, G. – CROVETTO, G. M. – FONDEVILA, M. – KRISHNAMOORTHY, U. – SINGH, B. – SPANGHERO, M. – STEINGASS, H. – ROBINSON, P. H. – KAILAS, M. M. 2002. Laboratory variation of 24 h *in vitro* production and estimated metabolizable energy values of ruminant feeds. *Animal Feed Science and Technology*, vol. 102, 2002, p. 169–180.
- GETACHEW, G. – MAKKAR, H. P. S. – BECKER, K. 2000. Stoichiometric relationship between short chain fatty acid and *in vitro* gas production in presence and absence of polyethylene glycol for tannin containing browses. In: EAAP Satellite Symposium, Gas Production: Fermentation Kinetics for Feed Evaluation and to Assess Microbial Activity, Wageningen, August 18-19, pp. 46–47.
- GIDAMIS, A. B. – PANGA, J. T. – SARWATT, S. V. – CHOVE, B. E. – SHAYO, N. B. 2003. Nutrients and antinutrient contents in raw and cooked leaves and mature pods of *Moringa oleifera*, Lam. *Ecology of Food and Nutrition*, vol. 42, 2003, p. 399–411.
- KOLJAJIC, V. – DORDEVIC, N. – GRUBIC, G. 1998. Effects of inoculants on ensiling of maize plant and alfalfa at different ratios. *Review of Research Work at the Faculty of Agriculture, Belgrade*, vol. 43 (2), 1998, p. 95–101.
- KUNG, L. – SHAVER, R. 2001: Interpretation and use of silage fermentation analysis reports. Forage Research Centre, Madison, Wisconsin, USA.
- LARBI, A. – SMITH, J. W. – ADEKUNLE, I. O. – KURDI, I. O. 1996. Studies on multipurpose fodder trees and shrubs in West Africa: Variation in determinants of forage quality in *Albizia* and *Paraserianthes* species. *Agroforestry Systems*, vol. 33, 1996, p. 1–11.
- LARBI, A. – SMITH, J. W. – RAJI, A. M. – KURDI, I. O. – ADEKUNLE, I. O. – LADIPO, D. O. 1997. Seasonal dynamics in dry matter degradation of browse in cattle, sheep and goats. *Small Ruminant Research*, vol. 25, 1997, p. 129–140.
- MAKKAR, H. P. S. – BECKER, K. 1997. Nutrients and antiquality factor in different morphological parts of the *Moringa oleifera* Tree. *Journal of Agricultural Science*, vol. 128, 1997, p. 311–322.
- MAN, N. V. – WIKTORSSON, H. 2002. Effect of molasses on nutritional quality of cassava and gliricidia tops silage. *Asian-Australasian Journal of Animal Science*, vol. 15, 2002, p. 1294–1299.
- MARSCHNER, H. 1993. Mineral Nutrition of Higher plant Second Edition. Academic press Ltd. London, Great Britain.
- MCDOWELL, L. R. – CONRAD, J. H. – HUMBRY, F. G. 1993. Minerals for grazing ruminants in tropical regions. Bulletin: The US Agency for Literature Development and Caribbean Basin Adversary Group (CBAG) USA.
- MCMENIMAN, N. P. – LITTLE, D. A. 1974. Studies on the supplementary feeding of sheep consuming mulga (*Acacia anura*). The provision of phosphorus and molasses supplement under grazing condition. *Australia Journal of Experimental Agriculture and Animal Husbandry*, vol. 14, 1974, p. 316–321.
- MENESES, M. D. – MEGIAS, J. – MADRID, A. – MARTINEZ-TERUEL, F. – HERNANDEZ, J. – OLIVA, J. 2007. Evaluation of the phytosanitary, fermentative and nutritive characteristics of the silage made from crude artichoke (*Cynara scolymus* L.) by-product feeding for ruminants. *Small Ruminant Research*, vol. 70, 2007, p. 292–296.
- MENKE, K. H. – STEINGASS, H. 1988. Estimation of the energetic feed value obtained from chemical analysis and gas production using rumen fluid. *Animal Research Development*, vol. 28, 1988, p. 7–55.
- MHERE, O. – MAASDORP, B. V. – TITTERTON, M. 1999. The effect of intercropping on production and ensilage of sweet forage sorghum and pennisetum with cowpea and dolichos bean. In: (Eds. T. Smith and W. Richards) Proceedings of Livestock Production Programme Workshop on Livestock Research in Zimbabwe. Department for International Development. Matopos, Zimbabwe.
- MUHAMMAD, I. R. – ABDULLAHI, B. – MOHAMMED, A. K. – TANKO, R. J. – KALLAH, M. S. – ALAWA, J. P. 2005. Influence of irrigation intervals on dry matter yield, concentration of crude protein, calcium and phosphorus in *Lablab purpureus* and *Sorghum alnum* fodder in the Sudan savannah zone of Nigeria. *Nigerian Journal of Animal Production*, vol. 32, 2005, p. 280–286.
- NASRULLAH, R. – NIIMI, M. – AKASHI, R. – KAWAMURA, O. 2004. Nutritive evaluation of forage plants grown in South Sulawesi, Indonesia. II. Mineral composition. *Asian-Australasian Journal of Animal Science*, vol. 17 (1), 2004, p. 63–67.
- NAYIGUHUGU, V. – KELLOGG, D. W. – LONGER, D. E. – JOHNSON, Z. B. 2002. Case study; performance and ensiling characteristics of tall-growing soybean lines used for silage. *American Registry of Professional Animal Scientist*, vol. 18 (1), 2002, p. 85–91.
- NORTON, B. W. 1994. Tree legumes as dietary supplements for ruminants. In: Gutteridge R C and Shelton H M (ed.). Forage tree legumes in tropical agriculture. CABI International, p. 192–201.
- OGUNBOSOYE, D. O. – BABAYEMI, O. J. 2010. Potential values of some non-leguminous browse plants as dry season feed for ruminants in Nigeria. *African Journal of Biotechnology*, vol. 9 (18), 2010, p. 2720–2726. Available online at <http://www.academicjournals.org/AJB>.

- OKOLI, I. C. – ANUNOBI, M. O. – OBUA, B. E. – ENEMUO, V. 2003. Studies on selected browses of southeastern Nigeria with particular reference to their proximate and some endogenous anti-nutritional constituents. *Livestock Research for Rural Development*, vol. 15 (9), 2003, <http://www.lrrd.org/lrrd15/9/okol159.htm>.
- PHIRI, M. S. – NGONGONI, N. T. – MAASDORP, B. V. – TITTERTON, M. – MUPANGWA, J. F. – SEBATA, A. 2007. Ensiling characteristics and feeding value of silage made from browse tree legume-maize mixtures. *Tropical and Subtropical Agroecosystems*, vol. 7, 2007, p. 149–156.
- ROGERS, P. – MURPHY, W. 2000. Levels of Dry Matter, Major Elements (calcium, magnesium, nitrogen, phosphorus, potassium, sodium and sulphur) and Trace Elements (cobalt, copper, iodine, manganese, molybdenum, selenium and zinc) in Irish Grass, Silage and Hay.
- SAS 2002. Statistical Analysis Systems user's guide (Statistics, Version. 8). SAS Institute Inc, Cary.
- UNDERWOOD, E. J. 1981. The mineral nutrition of livestock. 2nd edition. Commonwealth Agricultural Bureau, London.
- VAN SOEST, P. J. 1994. Nutritional Ecology of the Ruminant, 2nd edition. Comstock Publishing Associates/Cornell University Press, Ithaca, NY, USA.
- VAN SOEST, P. J. – ROBERTSON, J. B. – LEWIS, B. A. 1991. Methods for dietary fiber, neutral detergent fiber, and non-starch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*, vol. 74, 1991, p. 3583–3597.
- WATTIAUX, M. 1999. Introduction to silage-making. Dairy update. The Babcock Institute. University of Wisconsin. Feeding 502, 1999, p. 1–12, <http://babcock.cals.wisc.edu>.
- YOKOTA, H. – OHSHIMA, M. – HUANG, K. J. – OKAJIMA, T. 1995. Lactic acid production in Napier grass (*Pennisetum purpureum Schum.*) silage. *Grassland Science*, vol. 41, 1995, p. 207–211.
- ZANINE, A. – SANTOS, E. – FERREIRA, D. – OLIVEIRA, J. – ALMEIDA, J. – PEREIRA, O. 2006. Evaluation of Silage of Elephant-grass with addition of wheat meal. *Archives de Zootechnie*, vol. 55, 2006, p. 75–84.

Review

ETIOLOGY OF MASTITIS IN EWES AND POSSIBLE GENETIC AND EPIGENETIC FACTORS INVOLVED

V. TANČIN^{1,2*}, M. BAUER², I. HOLKO³, Š. BARANOVICH¹

¹Slovak University of Agriculture, FAFR, Department of Veterinary Science, Nitra, Slovak Republic

²NPPC – Research Institute for Animal Production Nitra, Slovak Republic

³VETSERVIS, s.r.o., Nitra, Slovak Republic

ABSTRACT

The paper reviews the knowledge related to the possible relationship of somatic cell counts to udder health and milk yield in dairy ewes. It attempts to point out the epigenetic and genetic aspects of udder health, methods for diagnostic of mastitis and pathogens involved. The possible physiological level of somatic cell counts (SCC) to establish the limits for ewe's milk are discussed. Data from the worldwide research are compared with limited results obtained in Slovakian dairy practice. Globally, applied research has focused on understanding of the relationship between SCC and mammary gland health through the presence of microorganisms. Many milk samples with high SCC are microbiologically negative, underlining the importance of research at the molecular level. Limits for SCC indicating health problems of ewe's udder are not yet established but during the last decades the proposal for limits decreased with time. Most paper considered 0.5×10^6 cells.mL⁻¹ or below for healthy udders. Data obtained from bulk milk showed that only 7.3 % of the samples were in the category below 0.5×10^6 cells.mL⁻¹ under Slovakian conditions. Possible genetic and epigenetic factors are discussed in this paper in relation to SCC. The identification of a genetic marker(s) that allows the inclusion of mastitis resistance in selection programmes would help to reduce the economic impact due to this disease. Subclinical mastitis is considered as a limiting factor for milk production. Several works have been published which presented a negative correlation between SCC and milk production.

Key words: mastitis; somatic cell counts; ewes; genetics; epigenetics; pathogens

INTRODUCTION

Sheep milk production is currently the main breeding aim of many agricultural farms and privately-employed farmers in many countries. Milk plays a crucial role in the economy of cooperatives and farms. The price of milk is affected, not only in the current situation, by the sheep milk market, but it is significantly influenced by the breeders. On the farmer's side, there are legislative limits for the total number of microorganisms in the milk delivered, which cannot exceed. Equally

important contributions of breeders are also hygienic safety of the milk for the consumers, especially in the marketing of milk and milk products directly sold on the farm. Some microorganisms found in milk may be a source of human health hazard (zoonoses), for example, *Staphylococcal* enterotoxins (Holečková *et al.*, 2004) which are not completely degraded in milk and milk products after pasteurization (Asao *et al.*, 2003). Total microorganisms in the delivered raw milk indicate the overall milking hygiene level and pathogenic species present in the milk thus representing the udder

*Correspondence: E-mail: vladimir.tancin@uniag.sk
Vladimír Tančin, Slovak University of Agriculture
FAFR, Department of Veterinary Science, Tr. A. Hlinku 2
949 76 Nitra, Slovak Republic

Received: December 7, 2015
Accepted: June 14, 2016

health status (Philpot and Nickerson, 1991). Moreover, microorganisms and udder health status adversely affect the technological qualities of the milk in its further processing (Leitner *et al.*, 2004). Recently, improvement in the overall immune responsiveness of the dairy cow including mammary gland against mastitis through the genetic and epigenetic factors has been believed to play a crucial role in providing better disease resistance, increasing animal welfare and food quality while maintain favourable production level to feed a growing population (Thompson-Crispi *et al.*, 2014).

The aim of this paper was to review the knowledge related to the possible relationship of somatic cell counts (SCC) to udder health and milk yield in dairy ewes. This work also attempts to point out the epigenetic and genetic aspects of udder health, methods for diagnosis of mastitis and pathogens involved.

Udder health status

The microorganisms present in the mammary gland represent one of the most serious diseases in dairy animals, called mastitis. The reaction of the body of ewes to the presence of microorganisms in the mammary gland is the transfer of blood white blood cells (somatic cells) into the milk. Diagnosis of mastitis is carried out by various methods, which differ in reliability, cost and complexity. The most common methods for mastitis detection in ewes are assessed on the basis of setting the number of somatic cells (SCC), bacteria culturing (McDougall *et al.*, 2001; Contreras *et al.*, 2007; Fragkou *et al.*, 2014) and palpation of the udder (Marogna *et al.*, 2010). Currently, real-time PCR is going to be intensively used for the diagnosis of mastitis pathogens (Zadoks *et al.*, 2014). More than 30 % of the samples from clinical and subclinical mastitis in dairy cows appear negative for the identification of pathogens during culturing (no growth of microorganisms). This situation is problematic for all stakeholders such as laboratories, farmers and veterinarians, which highlight the need for real-time PCR methods (Taponen *et al.*, 2009).

In addition to total SCC, various researchers have taken into consideration the presence of various types of white blood cells in the milk that may be related to the type of microorganisms in the udder, which allows to further specify the response of the body to the presence of various bacterial species (Ariznabarreta *et al.*, 2002; Bagnicka *et al.*, 2011; Leitner *et al.*, 2012). The knowledge of the art of mastitis in dairy ewes is in high demand at present because it is clearly confirmed that „sheep are not small cows“ (Zadoks *et al.*, 2014), because many pathogens are different from those that cause udder disease in dairy cows (Cuccuru *et al.*, 2011; Gilchrist *et al.*, 2013).

Physiological level of somatic cells in milk

At present in our country and across the world, individual as well as bulk samples of sheep milk are normally not analyzed for the presence of somatic cells, because the payment for the milk according to SCC is not implemented. This is due to the lack of objectively clarified factors and relationships that affect milk SCC in terms of physiological and pathological aspects (Fragkou *et al.*, 2014) despite the known fact that SCC is related to the presence of microorganisms in the mammary gland (McDougall *et al.*, 2002; Suarez *et al.*, 2002). In addition to the above mentioned reason, the determination of SCC is expensive for breeders of ewes (Adrias *et al.*, 2012).

Across the world, applied research is focused on understanding of the relationship between SCC and mammary gland health through the presence of microorganisms. Many milk samples with high SCC are microbiologically negative, underlining the importance of research at the molecular level (Zadoks *et al.*, 2014). At the physiological and pathophysiological level the SCC limit ranged from 0.25 to 1.0×10^6 cells.mL⁻¹ in the 80s, and the SCC for healthy udders was proposed at 0.5×10^6 cells.mL⁻¹ (Gonzalo and Gaudioso Lacasa, 1985). In a later work, Berthelot *et al.* (2006) reported healthy ewes with SCC below 0.5×10^6 cells and infected udders with SCC greater than 1×10^6 cells.mL⁻¹, while at the herd level, if SCC exceeded 0.65×10^6 cells.mL⁻¹, they indicated up to 15 % occurrence of mastitis. In determining relationship to milk production, Arias *et al.* (2012) recommended the limit value of 0.3×10^6 cells.mL⁻¹. In our work with 2,632 milk samples at the Experimental station, we have observed an increase of the proportion of ewes with SCC below 0.1×10^6 cells.mL⁻¹ from 31 % in 2010 to 56 % in 2013, and a decline in proportion of ewes with SCC less than 1×10^6 cells.mL⁻¹ from 21 % in 2010 to 12.5 % in 2013 (Idriss *et al.*, 2015). Recently in Tsigai ewes under practical conditions only 13 % of ewes had over 0.6×10^6 cells.mL⁻¹ (Vršková *et al.*, 2015). Another study from our experimental farm reported an average of 0.45×10^6 cells.mL⁻¹ with high variation coefficient (Margetin *et al.*, 2005, 2013). Earlier results published by Margetin *et al.* (1995; 1996) depending on year of study were 0.364×10^6 cells.mL⁻¹ (1993) and 1.1×10^6 cells.mL⁻¹ (1994) SCC during the period of suckling and milking but the data were obtained only from less than 50 animals per year. Available data from Slovakia indicate relatively good udder health of ewes but represent only very limited numbers of animals and farms. Riggio *et al.* (2013) stated that in uninfected Valle del Belice ewes, 83.7 % of the samples were in the category below 0.5×10^6 cells.mL⁻¹ and only 2.6 % above 1×10^6 cells.mL⁻¹. Similarly, a high percentage of samples with bacteriologically negative milk were in the following categories of SCC:

64.5 % had less than 0.05×10^6 cells.mL⁻¹, 81.9 % had less than 0.25×10^6 cells.mL⁻¹, and 92.4 % had less than 0.5×10^6 cells.mL⁻¹ (Pengov, 2001). Therefore the author considered the threshold of 0.25×10^6 cells.mL⁻¹ beyond the assessment of udder health.

In spite of the very good udder health in selected farms reported by Idriss *et al.* (2015) and Vršková *et al.* (2015) the health of udder in Slovakian sheep farms seems to be much more complicated and difficult. The bulk milk analysis of 1086 samples from March to August revealed that only 7.3 % of the samples were in the category below 0.5×10^6 cells.mL⁻¹ (Tomaška *et al.*, 2015). In last mentioned report 49 % of samples were over 1×10^6 cells.mL⁻¹ which deserves more detailed research in mastitis control in dairy practice in Slovakia. In general Slovakian breeders do not test for SCC in ewes' milk, and have no information about the health status of the mammary glands and thus the potential risks of the sale of raw unpasteurized sheep milk and sheep milk products to consumers' health. Farmers lack information on risks of breeding environment for udder health, including economic losses caused by mastitis. Even though SCC as an indicator of udder health and milk quality is not part of pricing milk, countries with high level of sheep breeding pay considerable attention to udder health (Leitner *et al.*, 2012; Fragkou *et al.*, 2014). Research in these countries is aimed to improve the health of the animal's mammary gland, resulting in improved economy of breeding ewes, the competitiveness of enterprises and production of hygienically and nutritionally better milk quality. Slovakia is also considered as the country with well-developed traditional sheep breeding, which requires more intensive research in the production of safe and hygienic raw sheep milk in primary production. Such research will also contribute to a more effective breeding of dairy ewes.

Classification of mastitis

In terms of the most common occurrence of mastitis, in practice, this disease can be divided into clinical and subclinical mastitis. Clinical mastitis is quickly detected by the breeders on the basis of clinical symptoms, such as painful udder, edema and changes in the consistency of the milk (Marogna *et al.*, 2010), where the producer may use such symptoms for disposal of such milk from delivering into the dairy and in terms of the risk of microbial contamination, also from the food chain. More detrimental to the sheep breeding are subclinical mastitis, which do not show any clinical signs at the udder and milk level. Subclinical mastitis negatively influences cheese yield and its quality (Silanikove *et al.*, 2014). Moreover, such milk is not only a source of pathogens but also their byproducts - toxins (Asao *et al.*, 2003), which have been confirmed

in our conditions (Holečková *et al.*, 2004; Mašlanková *et al.*, 2009; Zigo *et al.*, 2014). The finding that the subclinical mastitis is the most widespread in the breeding system of ewes is also important (Bergonier *et al.*, 2003; Leitner *et al.*, 2004). In farms with a good management, the incidence of clinical mastitis is below 5 % (McDougall *et al.*, 2001), but subclinical mastitis have been detected at 15-40 % of the ewes (Kiossis, *et al.*, 2007, Contreras *et al.*, 2007).

Mastitis pathogens

As indicated above, subclinical mastitis is a limiting factor for milk production. Several works have been published which present a negative correlation between SCC and milk production (Fuentes *et al.*, 1998, Gonzalo *et al.*, 2002). Major pathogens (infectious; *Staphylococcus aureus*, *Streptococcus agalactiae*, *Mycoplasma*) cause a significant increase in SCC and decline in milk production compared to minor microorganisms (Gonzalo *et al.*, 2002; Riggio *et al.*, 2013). On the other hand, minor microorganisms (environmental; the most common coagulase negative staphylococci - CNS) are the predominant organisms in the milk of ewes in the herd (Gonzalo *et al.*, 2002; Kiossis *et al.*, 2007; Pilipčincová *et al.*, 2010; Marogna *et al.*, 2010, Bagnicka *et al.*, 2011) and deserve deeper attention (Leitner *et al.*, 2012). In farms of eastern Slovakia, Pilipčincová *et al.* (2010) analyzed the presence of CNS species in individual ewe's milk from farms with mostly hand-milking (94.5 %), which differed significantly from the species present in cow milk (Zadoks and Watts, 2009). Specification of CNS species increases the effectiveness of different control treatments against mastitis (Ruegg, 2009). Significant differences in detection of different CNS species between regions in Slovakia (Pilipčincová *et al.*, 2010) and in other countries (Mavrogianni *et al.*, 2007), as well as the specific presence of microorganisms in sheep milk versus cow milk (Zadoks *et al.*, 2009) suggest the need for a deeper analysis of mastitis pathogens in ewes in Slovakia and the identification of risk factors of external and internal environment. Microscopic fungi and yeasts play a certain role in the infection of the udder of dairy cow (Scaccabarozzi *et al.*, 2011; Idriss *et al.*, 2013), which draw little scientific attention in sheep milk.

Mastitis and milk yield

The health status of the mammary gland, manifested by increased SCC is under considerable attention of researchers also in relation to the drop in milk production. Negative phenotypic correlation between SCC and milk production in different breeds was reported by several authors in Manchega ewes (Adrias *et al.*, 2012) and in Churra ewes (Gonzalo *et al.*,

2002). Špánik *et al.* (1996) calculated the negative correlation between SCC and the yield in the experimental herd. Negative significant correlation between SCC and milk production in Tsigai ewes during both suckling and milking period was also found out by Margetin *et al.* (1996). Recently we have reported non-significant reduction of milk yield with increasing SCC in Tsigai ewes (Vršková *et al.*, 2015). The differences in performance between ewes with SCC above and below 0.3×10^6 cells.mL⁻¹ increased during lactation (Adrias *et al.*, 2012). The authors also found that ewes with twins had a higher proportion of samples with SCC above 0.3×10^6 cells.mL⁻¹, but also higher milk production than ewes with one lamb (Adrias *et al.*, 2012). Similarly, Olives *et al.* (2013) noted a lower milk production of infected udder by 16 % during 7 weeks of lactation, and in comparing the two halves of the udder, healthy udder-half compensated for the loss of the second half by up to 6.6 %. Similar reduction of milk yield between two quarters in the same front/rear position with different SCC was also reported in cows (Tančin and Uhrinčat', 2014).

Possible genetic markers

The identification of a genetic marker(s) that allows for the inclusion of mastitis resistance in selection programmes would help reduce the economic impact due to this disease. However, the selection of a candidate gene is a difficult task because mastitis is a complex disease influenced by many genes and environmental factors. Problems associated with breeding directly for mastitis resistance also include its low heritability, ranging from 0.02-0.10 (Nash *et al.*, 2000). With the recent development of extensive high-throughput genomic tools research efforts have increasingly turned to identifying single nucleotide polymorphisms (SNPs) associated with resistance as well as quantifying the genetic control of the host-pathogen interaction (Rupp and Foucras, 2010). The solutions to improve resistance to mastitis are likely to be those that focus on information from genome – wide association studies (GWAS), or selection based on breeding values of immune responses, which take into account complex genetic interactions between the innate and adaptive host defense mechanisms without the necessity of knowing all about each individual gene. These approaches may be best suited to help alleviate mastitis, at least until we gain more knowledge about genetic and epigenetic regulation of host defense mechanisms.

Major histocompatibility complex (MHC) plays an important role in immunological defense against pathogens. MHC is a group of genes on a single chromosome that codes the MHC antigens and includes two major subfamilies: class I and class II genes. MHC molecules that function in the recognition

event, which is termed “antigen presentation”, are polymorphic glycoproteins found on cell membranes. The MHC participates in the development of both humoral and cell mediated immune responses. The general consensus is that these genes may be useful as genetic markers of a higher or lower risk of mastitis in cows (Conington *et al.*, 2008).

In sheep, MHC is located on chromosome 20 and is called Ovar (Hediger *et al.*, 1991). A complete ovine MHC sequence map was assembled by successful shotgun sequencing of 26 overlapping BAC clones (Gao *et al.*, 2010). The MHC of sheep and cattle share orthologous class II DR and DQ A and B loci with rodents and primates. A single dominant and highly polymorphic DRB locus encoding the beta chain of the MHC class II DR heterodimer has been described in domestic sheep Ovar-DRB1 (Scott *et al.*, 1991). The detailed genomic organization and allelic diversity of Ovar-DRB1 locus has been described (Ballingall *et al.*, 2008). Recently, unusual allelic diversity has also been identified at the DRA locus in domestic sheep (Ballingall *et al.*, 2010). Several studies have shown the existence of ovine class II loci that are homologous to HLA-DQB (Dukkipati *et al.*, 2006). As in other vertebrate species, a high degree of polymorphism is found in the Ovar-DQB genes, with most of the polymorphic sites located in exon 2, which encodes for the antigen-binding site.

It has been reported that alleles of different MHC genes correlate with disease resistance in sheep. Currently, relevant research on Ovar genes' polymorphism and disease resistance or susceptibility mainly concentrates on Ovar-DRB1 and Ovar-DQB. Herrmann-Hoesing *et al.* (2008) detected that Ovar-DRB1 alleles contribute as a host genetic factor controlling ovine pneumonia provirus levels. Larruskain *et al.* (2010) found a significant association of Ovar-DRB1 alleles with resistance to Maedi-Visna and pulmonary adenocarcinoma viruses. Recently, Shen *et al.* (2014) have shown that polymorphism in Ovar-DRB1/DQB1 can be used as a marker of resistance to Echinococcosis in Chinese Merino sheep.

Epigenetic factors

Udder disease is affected by a number of external environmental factors (management, manner of milk removal and milking technology, season) and internal factors (physiological status of the body, like stage and order of lactation, oestrus, udder shape and response of ewes to milking, condition) (Raynal-Ljutovac *et al.*, 2007). Other important factors include the overall level of husbandry and management aimed to reduce the risk of mastitis, such as drying ewes that are treated with antibiotics (Shwimmer *et al.*, 2008; Spanu *et al.*, 2011) and in particular the routine of machine milking (Leitner *et al.*, 2008). The milking routine must be based

on biological needs of ewes for machine milking and both from the morphological as well as physiological aspects (Mačuhová *et al.*, 2008, 2012; Tančin *et al.*, 2011; Antonič *et al.*, 2013b). Our results from several farms showed that ewes bred in Slovakia have a relatively poor response to machine milking, which is observed as retention of milk in the udder. Holding of the milk in the udder reduces milk production (Silanikove *et al.*, 2010) and increases the risk of disease of the udder. It is very important to further specify the state of health of the mammary gland of ewes bred in Slovakian conditions. Furthermore, to assess the impact of manner of milk removal (hand, machine) on udder health in relation to breed, functionality and parameters of milking equipment, ewes age, stage of lactation, milking frequency, number of lambs, organization of work in the milking process and the overall level of farming must be studied. We earlier reported an increased risk for udder health-related to weaning lambs and ewes adaptation to machine milking (Antonič *et al.*, 2013a), and changes in conditions of milking (Kulinová *et al.*, 2012; Jackuliaková *et al.*, 2014; Tančin *et al.*, 2015). It seems that hand milking induces more udder infection than machine milking and the type of milking equipment (mobile and stationary) plays a role in the risk of mastitis as well (Marogna *et al.*, 2010). More frequent health problems of udder for milking with mobile devices compared with stationary ones have been noted by Marogna *et al.* (2010), which justify the worse parameters of mobile milking equipment (vacuum stability, frequency pulsations, activity of liner). The transition from extensive to intensive dairy sheep farming system also brings more infections caused by environmental pathogens (Marogna *et al.*, 2010). In Slovakian conditions these were not studied though many changes have taken place in milk removal systems in dairy practice in Slovakia.

The completeness of udder emptying and speed of milking of cows are considered among the most important indicators of the impact of technology on milk removal and handling of dairy animals (Tančin and Bruckmaier, 2001), which demonstrate not only the welfare of the animals during milking, but are one of the important reactions of the organism in reducing the risk of udder disease (Tančin *et al.*, 2006, 2007). Although ewes have certain differences in milk distribution in the udder as compared to dairy cows, it is likely that the speed and completeness of milking to keep good udder health is equally important. Thus, intense transition from hand to machine milking of ewes under practical conditions of Slovakia deserves serious research on the prevention of udder diseases.

CONCLUSIONS

From the above mentioned information it could be concluded that without the knowledge related to udder health and possible risk factors of breeding systems in dairy practice in Slovakia, it would not be possible to further improve the development of sheep breeding in the country, where sheep are considered to be of high importance for animal production, sustainability of countryside and production of special milk products.

ACKNOWLEDGEMENTS

This paper was prepared as a part of realization of the project Kega 006SPU-4/2014 "Modernization of teaching of domestic animal hygiene" and the project „MLIEKO no. 26220220196“ supported by the Operational Programme for Research and Development from the European Regional Development Fund.

REFERENCES

- ANTONIČ, J. – MAČUHOVÁ, L. – UHRINČAŤ, M. – TANČIN, V. 2013a. The effect of milk ejection occurrence before or during machine milking on milkability and milk composition of ewes. *Veterinaria ir Zootechnika (Vet Med Zoot)*, vol. T. 61 (83), 2013a, p. 3–7.
- ANTONIČ, J. – TANČIN, V. – UHRINČAŤ, M. – MAČUHOVÁ, L. – MAČUHOVÁ, J. – JACKULIAKOVÁ, L. 2013b. The effect of exogenous oxytocin on milkability and milk composition in ewes differed in milk flow pattern. *Small Ruminant Research*, vol. 113, 2013b, p. 254–257.
- ARIAS, R. – OLLETE, B. – RAMON, M. – ARIAS, C. – GALLEGU, R. – MONTORO, V. – GONZALO, C. – PEREZ-GUZMAN, M. D. 2012. Long-term study of environmental effects on test-day somatic cell count and milk yield in Manchega sheep. *Small Ruminant Research*, vol. 106, 2012, p. 92–97.
- ARIZNABARRETA, A. – GONZALO, C. – SAN PRIMITIVO, F. 2002. Microbiological quality and somatic cell count of ewes milk with special reference to staphylococci. *Journal of Dairy Science*, vol. 85, 2002, p. 1370–1375.
- ASAO, T. – KUMEDA, Y. – KAWAI, T. – SHIBATA, T. – ODA, H. – HARUKI, K. – NAKAZAWA, H. – KOZAKI, S. 2003. An extensive outbreak of staphylococcal food poisoning due to low-fat milk in Japan: estimation of enterotoxin A in the incriminated

- milk and powdered skim milk. *Epidemiology and Infection*, vol. 130, 2003, p. 33–40.
- BAGNICKA, E. – WINNICKA, A. – JOZWIK, A. – RZEWUSKA, M. – STRZALKOWSKA, N. – KOSCIUCZUK, E. – PRUSAK, B. – KABA, J. – HORBANCZUK, J. – KRZYZEWSKI, J. 2011. Relationship between somatic cell count and bacterial pathogens in goat milk. *Small Ruminant Research*, vol. 100, 2011, p. 72–77.
- BALLINGALL, K. T. – FARDOE, K. – MCKEEVER, D. J. 2008. Genomic organisation and allelic diversity within coding and non-coding regions of the Ovar-DRB1 locus. *Immunogenetics*, vol. 60 (2), 2008, p. 95–103.
- BALLINGALL, K. T. – ROCCHI, M. S. – MCKEEVER, D. J. – WRIGHT, F. 2010. Trans-Species Polymorphism and Selection in the MHC Class II DRA Genes of Domestic Sheep. *PLoS ONE*, vol. 5 (6), 2010, p. 1–10.
- BERGONIER, D. – DE CREMOUX, R. – RUPP, R. – LAGRIFFOUL, G. – BERTHELOT, X. 2003. Mastitis in dairy small ruminants. *Veterinary Research*, vol. 34, 2003, p. 689–716.
- BERTHELOT, X. – LAGRIFFOUL, G. – CONCORDET, D. – BARILLET, F. – BERGONIER, D. 2006. Physiological and pathological thresholds of somatic cell counts in ewe milk. *Small Ruminant Research*, vol. 62, 2006, p. 27–31.
- CONINGTON, J. – CAO, G. – STOTT, A. – BÜNGER, L. 2008. Breeding for resistance to mastitis in United Kingdom sheep, a review and economic appraisal. *The Veterinary Record*, vol. 162 (12), 2008, p. 369–376.
- CONTRERAS, A. – SIERRA, D. – SANCHEZ, A. – CORRALES, J. C. – MARCO, J. C. – PAAPE, M. J. – GONZALO, C. 2007. Mastitis in small ruminants. *Small Ruminant Research*, vol. 68, 2007, p. 145–153.
- CUCCURU, C. – MELONI, M. – SALA, E. – SCACCABAROZZI, L. – LOCATELLI, C. – MORONI, P. – BRONZO, V. 2011. Effect of intramammary infection on somatic cell score and milk yield in Sarda sheep. *New Zealand Veterinary Journal*, vol. 59, 2011, p. 128–131.
- DUKKIPATI, V. S. R. – BLAIR, H. T. – GARRICK, D. J. – MURRAY, A. 2006. Ovar-Mhc' – ovine major histocompatibility complex: structure and gene polymorphisms. *Genetics and Molecular Research*, vol. 5 (4), 2006, p. 581–608.
- FRAGKOU, I. A. – BOSCOS, C. M. – FTHENAKIS, G. C. 2014. Diagnosis of clinical or subclinical mastitis in ewes. *Small Ruminant Research*, vol. 118, 2014, p. 86–92.
- FUERTES, J. A. – GONZALO, C. – CERRIEDO, J. A. – SAN PRIMITIVO, F. 1998. Parameters of test day milk yield and milk components in dairy ewes. *Journal of Dairy Science*, vol. 81, 1998, p. 1300–1307.
- GAO, J. – LIU, K. – LIU, H. – BLAIR, T. – LI, G. – CHEN, CH. – TAN, P. – MA, R. Z. 2010. A complete DNA sequence map of the ovine Major Histocompatibility Complex. *BMC Genomics*, vol. 11 (1), 2010, p. 466.
- GILCHRIST, T. L. – SMITH, D. G. – FITZPATRICK, J. L. – ZADOKS, R. N. – FONTAINE, C. 2013. Comparativemolecular analysis of ovine and bovine Streptococcus uberis isolates. *Journal of Dairy Science*, vol. 96, 2013, p. 962–970.
- GONZALO, C. – GAUDIOSO LACASA, V. R. 1985. Evolution des types cellulaires du laits de bredis (race Churra) en fonction des dénombrements cellulaires totaux pendant latraite mécanique et manuelle. *Annales Zootechnie*, vol. 34, 1985, p. 257–264.
- GONZALO, C. – ARIZNABARRETA, A. – CARRIEDO, J. A. – SAN PRIMITIVO, F. 2002. Mammary pathogens and their relationship to somatic cell count and milk yield losses in dairy ewes. *Journal of Dairy Science*, vol. 85, 2002, p. 1460–1467.
- HEDIGER, R. – ANSARI, H. A. – STRANZINGER, G. F. 1991. Chromosome banding and gene localizations support extensive conservation of chromosome structure between cattle and sheep. *Cytogenetics and Cell Genetics*, vol. 57 (2-3), 1991, p. 127–134.
- HERRMANN-HOESING, L. M. – WHITE, S. N. – MOUSEL, M. R. – LEWIS, G. S. – KNOWLES, D. P. 2008. Ovine progressive pneumonia provirus levels associate with breed and Ovar-DRB1. *Immunogenetics*, vol. 60 (12), 2008, p. 749–758.
- HOLEČKOVÁ, B. – KALIŇAČOVÁ, V. – GONDOL, J. – FOTTA, M. – HOLODA, E. – BELIČKOVÁ, E. 2004. Production of enterotoxins by *Staphylococcus aureus* isolated from sheep milk. *Bulletin of the Veterinary Institute in Pulawy*, vol. 48, 2004, p. 41–45.
- IDRISS, S. H. – TANČIN, V. – FOLTÝS, V. – KIRCHNEROVÁ, K. – TANČINOVÁ, D. – VRŠKOVÁ, M. 2013. Relationship between mastitis causative pothogens and somatic cell counts in milk of dairy cows. *Potravinárstvo*, vol. 7 (1), 2013, p. 207–212.
- IDRISS, S. E. – TANČIN, V. – MARGETÍN, M. – TANČINOVÁ, D. – SLÁMA, P. – HAVLÍČEK, Z. 2015. The frequency of distribution of somatic cell count in dairy ewe's milk. *Journal of Microbiology, Biotechnology and Food Sciences*, vol. 4, 2015, p. 148–151.
- JACKULIAKOVÁ, L. – TANČIN, V. – UHRINČAĎ, M. – MAČUHOVÁ, L. – ANTONIČ, J. – ORAVCOVÁ, M. – SLÁMA P. 2014. The effect of ewe's relocation on milk composition and milk flow kinetics. *Potravinárstvo*, vol. 8, 2014, p. 135–140.
- KIOSSIS, E. – BROZOS, C. N. – PETRIDOU, E. – BOSCOS, C. 2007. Program for the control of subclinical mastitis in dairy Chios breed ewes during

- lactation. *Small Ruminant Research*, vol. 73, 2007, p. 194–199.
- KULINOVÁ, K. – MAČUHOVÁ, L. – UHRINČAĎ, M. – TANČIN, V. 2012. The effect of stressful treatment before and during milking on milkability of dairy ewes. *Veterinarija ir Zootechnika (Vet Med Zoot)*, vol. T. 57 (79). 2012, p. 39–43.
- LARRUSKAIN, A. – MINGUIJÓN, E. – GARCÍA-ETXEBARRIA, K. – MORENO, B. – AROSTEGUI, I. – JUSTE, R. A. – JUGO, B. M. 2010. MHC class II DRB1 gene polymorphism in the pathogenesis of Maedi–Visna and pulmonary adenocarcinoma viral diseases in sheep. *Immunogenetics*, vol. 62 (2), 2010, p. 75–83.
- LEITNER, G. – CHAFFER, M. – SHAMAY, A. – SHAPIRO, F. – MERIN, U. – EZRA, E. – SARAN, A. – SILANIKOVE, N. 2004. Changes in Milk Composition as Affected by Subclinical Mastitis in Sheep. *Journal of Dairy Science*, vol. 87, 2004, p. 46–52.
- LEITNER, G. – MERIN, U. – KRIFUCK, O. – BLUM, S. – RIVAS, A. L. – SILANIKOVE, N. 2012. Effects of intra-mammary bacterial infection with coagulase negative staphylococci and stage of lactation on shedding of epithelial cells and infiltration of leukocytes into milk: Comparison among cows, goats and sheep. *Veterinary Immunology and Immunopathology*, vol. 147, 2012, p. 202–210.
- LEITNER, G. – SILANIKOVE, N. – MERIN, U. 2008. Estimate of milk and curd yield loss of sheep and goat with intramammary infection and its relation to somatic cell counts. *Small Ruminant Research*, vol. 74, 2008, p. 221–225.
- MAČUHOVÁ, L. – UHRINČAĎ, M. – MAČUHOVÁ, J. – MARGETIN, M. – TANČIN, V. 2008. The first observation of milkability of the sheep breeds Tsigai, Improved Valachian and their crosses with Lacaune. *Czech Journal of Animal Science*, vol. 53, 2008, p. 528–536.
- MAČUHOVÁ, L. – TANČIN, V. – UHRINČAĎ, M. – MAČUHOVÁ, J. 2012. The level of udder emptying and milk flow stability in Tsigai, Improved Valachian, and Lacaune ewes during machine milking. *Czech Journal of Animal Science*, vol. 57, 2012, p. 240–247.
- MARGETIN, M. – ČAPISTRÁK, A. – ŠPÁNIK, J. – FOLTYS, V. 1996. Somatic cells in sheep milk in relation to milk production and composition during sucking and milking. *Živočišná výroba*, vol. 41, 1996, p. 543–550.
- MARGETIN, M. – ČAPISTRÁK, A. – VALKOVSKY, P. – ŠPÁNIK, J. – FOLTYS, V. 1995. Variation in somatic-cell counts in ewes milk during lactation. *Živočišná výroba*, vol. 40, 1995, p. 257–261.
- MARGETIN, M. – MILERSKI, M. – APOLEN, D. – ČAPISTRÁK, A. – ORAVCOVÁ, M. – DEBRECENI, O. 2013. Relationships between production, quality of milk and udder health status of ewes during machine milking. *Journal of Central European Agriculture*, vol. 14, 2013, p. 328–340.
- MARGETIN, M. – ŠPÁNIK, J. – MILERSKI, M. – ČAPISTRÁK, A. – APOLEN, D. 2005. Relationships between morphological and functional udder traits and somatic cell count in milk of ewes. ICAR Technical Series No. 10, ed. Tančin, V., Mihina, S., Uhrinčáď, M., p. 255–258, ISBN 92-95014-07-3.
- MAROGNA, G. – ROLESU, S. – LOLLAI, S. – TOLA, S. – LEORI, G. 2010. Clinical findings in sheep farms affected by recurrent bacterial mastitis. *Small Ruminant Research*, vol. 88, 2010, p. 119–125.
- MAŠLANKOVÁ, J. – PILIPČINCOVÁ, I. – TKÁČIKOVÁ, E. 2009. Pheno- and genotyping of *Staphylococcus aureus* isolates of sheep origin. *Acta Veterinaria Brno*, vol. 78, 2009, p. 345–352.
- MAVROGIANNI, V. S. – CRIPPS, P. J. – FTHENAKIS, G. C. 2007. Bacterial flora and risk of infection of the ovine teat duct and mammary gland throughout lactation. *Preventive Veterinary Medicine*, vol. 79, 2007, p. 163–173.
- MCDUGALL, S. – MURDOUGH, P. A. – PANKEY, W. – DELANEY, C. – BARLOW, J. – SCRUTON, D. 2001. Relationships among somatic cell count, California mastitis test, impedance and bacteriological status of milk in goat and sheep in early lactation. *Small Ruminant Research*, vol. 40, 2001, p. 245–254.
- MCDUGALL, S. – PANKEY, W. – DELANEY, C. – BARLOW, J. – MURDOUGH, P. A. – SCRUTON, D. 2002. Prevalence and incidence of subclinical mastitis in goat and dairy ewes in Vermont, USA. *Small Ruminant Research*, vol. 46, 2002, p. 115–121.
- NASH, D. L. – ROGERS, G. W. – COOPER, J. B. – HARGROVE, G. L. – KEOWN, J. F. – HANSEN, L. B. 2000. Heritability of clinical mastitis incidence and relationships with sire transmitting abilities for somatic cell score, udder type traits, productive life and protein yield. *Journal of Dairy Science*, vol. 83 (10), 2000, p. 2350–2360.
- OLIVES DE, A. M. – DÍAZ, J. R. – MOLINA, M. P. – PERIS, C. 2013. Quantification of milk yield and composition changes as affected by subclinical mastitis during the current lactation in sheep. *Journal of Dairy Science*, vol. 96, 2013, p. 7698–7708.
- PENGOV, A. 2001. The role of coagulase-negative *Staphylococcus* spp. and associated somatic cell count in the ovine mammary gland. *Journal of Dairy Science*, vol. 84, 2001, p. 572–574.
- PHILPOT, N. – NICKERSON, S. C. 1991. Mastitis: counter attack. Babson Bros. Company, p. 149.
- PILIPČINCOVÁ, I. – BHIDE, M. – DUDŘÍKOVÁ, E. – TRÁVNÍČEK, M. 2010. Genotypic characterisation, of coagulase-negative staphylococci isolated from

- sheep milk in Slovakia. *Acta Veterinaria Brno*, vol. 79, 2010, p. 269–275.
- RAYNAL-LJUTOVAC, K. – PIRISI, A. – CREMOUX DE, R. – GONZALO, C. 2007. Somatic cells of goat and sheep milk: analytical, sanitary, productive and technological aspects. *Small Ruminant Research*, vol. 68, 2007, p. 126–144.
- RIGGIO, V. – PESCE, L. L. – MORREALE, S. – PORTOLANO, B. 2013. Receiver-operating characteristic curves for somatic cell scores and california mastitis test in Valle del Belice dairy sheep. *The Veterinary Journal*, vol. 196, 2013, p. 528–532.
- RUEGG, P. L. 2009. The quest for the perfect test: Phenotypic versus genotypic identification of coagulase-negative staphylococci associated with bovine mastitis. *Veterinary Microbiology*, vol. 134, 2009, p. 15–19.
- RUPP, R. – FOUCRAS, G. 2010. Genetics of Mastitis in Dairy Ruminants. In: *Breeding for Disease Resistance in Farm Animals*, eds. S. C. Bishop et al., 3rd Edition, CABI, 2010.
- SCACCABAROZZI, L. – LOCATELLI, C. – PISONI, G. – MANAROLLA, G. – CASULA, A. – BRONZO, V. – MORONI, P. 2011. Short communication: Epidemiology and genotyping of *Candida rugosa* strains responsible for persistent intramammary infections in dairy cow. *Journal of Dairy Science*, vol. 94, 2011, p. 4574–4577.
- SCOTT, P. C. – MADDUX, J. F. – GOGOLIN-EWENS, K. J. – BRANDON, M. R. 1991. The nucleotide sequence and evolution of ovine MHC class II B genes: *DQB* and *DRB*. *Immunogenetics*, vol. 34 (2), 1991, p. 80–87.
- SHWIMMER, A. – KENIGSWALD, G. – VAN STRATEN, M. – LAVI, Y. – MERIN, U. – WEISBLIT, L. – LEITNER, G. 2008. Dry-off treatment of Assaf sheep: Efficacy as a management tool for improving milk quantity and quality. *Small Ruminant Research*, vol. 74, 2008, p. 45–51.
- SHEN, H. – HAN, G. – JIA, B. – JIANG, S. – DU, Y. 2014. MHC-*DRB1/DQB1* gene polymorphism and its association with resistance/susceptibility to cystic Echinococcosis in Chinese Merino sheep. *Journal of Parasitology Research*, 2014, ID 272601, p. 7.
- SILANIKOVE, N. – LEITNER, G. – MERIN, U. – PROSSER, C. G. 2010. Recent advantages in exploiting goats milk: quality, safety and production aspects. *Small Ruminant Research*, vol. 89, 2010, p. 110–124.
- SILANIKOVE, N. – MERIN, U. – LEITNER, G. 2014. On effects of subclinical mastitis and stage of lactation on milk quality in goats. *Small Ruminant Research*, vol. 122, 2014, p. 76–82.
- SPANU, C. – BERGER, Y. M. – THOMAS, D. L. – RUEGG, P. L. 2011. Impact of intramammary antimicrobial dry treatment and teat sanitation on somatic cell count and intramammary infection in dairy ewes. *Small Ruminant Research*, vol. 97, 2011, p. 139–145.
- SUAREZ, V. H. – BUSETTI, M. R. – MIRANDA, A. O. – CALVINHO, L. F. – BEDOTTI, D. O. – CANAVESIO, V. R. 2002. Effect of infectious status and parity on somatic cell count and California mastitis test in pampinta dairy ewes. *Journal of Veterinary Medicine*, vol. 49 (5), 2002, p. 230–234.
- ŠPÁNIK, J. – KAČINCOVÁ, A. – MARGETÍN, M. – ČAPISTRÁK, A. – KALIŠ, M. 1996. Dependence of sheep milk quality on somatic cell counts. *Journal of Farm of Animal Science*, vol. 29, 1996, p. 111–116.
- TANČIN, V. – BRUCKMAIER, R. M. 2001. Factors affecting milk ejection and removal during milking and suckling of dairy cows. *Veterinary Medicine-Czech*, vol. 46, 2001, p. 108–118.
- TANČIN, V. – IPEMA, A. H. – HOGWERF P. 2007. Interaction of Somatic Cell Count and Quarter Milk Flow Patterns. *Journal of Dairy Science*, vol. 90, 2007, p. 2223–2228.
- TANČIN, V. – IPEMA, I. H. – HOGWERF, P. H. – MAČUHOVÁ, J. 2006. Sources of variation in milk flow characteristics at udder and quarter levels. *Journal of Dairy Science*, vol. 89, 2006, p. 978–988.
- TANČIN, V. – MAČUHOVÁ, J. – JACKULIAKOVÁ, L. – UHRINČAĎ, M. – ANTONIČ, J. – MAČUHOVÁ, L. – JÍLEK, F. 2015. The effect of social stress on milking efficiency in dairy ewes differed in milk flow kinetic. *Small Ruminant Research*, vol. 125, 2015, p. 115–119.
- TANČIN, V. – MAČUHOVÁ, L. – ORAVCOVÁ, M. – UHRINČAĎ, M. – KULINOVÁ, K. – ROYCHOUDHURY, S. – MARNET P. G. 2011. Milkability assessment of Tsigai, Improved Valachian, Lacaune and F1 Crossbred ewes (Tsigai×Lacaune, Improved Valachian×Lacaune) throughout lactation. *Small Ruminant Research*, vol. 97, 2011, p. 28–33.
- TANČIN, V. – UHRINČAĎ, M. 2014. The effect of somatic cell on milk yield and milk flow at quarter level. *Veterinarija ir Zootechnika (Vet Med Zoot)*, vol. 66, 2014, p. 69–72.
- TAPONEN, S. – SALMIKIVI, L. – SIMOJOKI, H. – KOSKINEN, M. T. – PYÖRÄLÄ, S. 2009. Real-time polymerase chain reaction-based identification of bacteria in milk samples from bovine clinical mastitis with no growth in conventional culturing. *Journal of Dairy Science*, vol. 92, 2009, p. 2610–2617.
- TOMÁŠKA, M. – HOFERICOVÁ, M. – KLIMEŠOVÁ, M. – HANUŠ, O. – VORLOVÁ, L. – KOLOŠTA, M. 2015. Occurrence of somatic cells in bulk samples of raw sheep's milk. *Zborník Hygiena a technologic potravín XLV. Lenfeldovy a Höklovy dny*, 2015, p. 197–200. ISBN: 978-80-7305-762-6.

-
- THOMPSON-CRISPI, K. – ATALLA, H. – MIGLIOR, F. – MALLARD, B. A. 2014. Bovine mastitis: frontiers in immunogenetics. *Frontier in Immunology*, vol. 5 (article 493), 2014, p. 1–10.
- VRŠKOVÁ, V. – TANČIN, V. – KIRCHNEROVÁ, K. – SLÁMA, P. 2015. Evaluation of daily milk production in tsigai ewes by somatic cell count. *Potravinarstvo*, vol. 9, 2015, p. 206–210.
- ZADOKS, R. N. – WATTS, J. L. 2009. Species identification of coagulase-negative staphylococci: genotyping is superior to phenotyping. *Veterinary Microbiology*, vol. 134, 2009, p. 20–28.
- ZADOKS, R. N. – TASSI, R. – MARTIN, E. – HOLOPAINEN, J. – McCALLUM, S. – GIBBONS, J. – BALLINGALL, K. T. 2014. Comparison of bacteriological culture and PCR for detection of bacteria in ovine milk – sheep are not small cows. *Journal of Dairy Science*, vol. 97, 2014, p. 6326–6333.
- ZIGO, F. – VASIL, M. – ELEČKO, J. – FARKAŠOVÁ, Z. – LAPIN, M. 2014. Production of enterotoxins of *Staphylococcus* spp. isolated from samples of sheep milk. *Potravinarstvo*, vol. 8, 2014, p. 92–96.
-

Short communication

THE POTENTIAL USE OF CHICKEN BLASTODERMAL CELLS OF ORAVKA BREED

A. SVORADOVÁ^{1*}, L. KUŽELOVÁ², P. CHRENEK^{3,4}

¹Faculty of Natural Sciences, Constantine the Philosopher University, Nitra, Slovak Republic

²The Research Centre AgroBioTech Nitra, Slovak Republic

³NPPC – Research Institute for Animal Production Nitra, Slovak Republic

⁴Faculty of Biotechnology and Food Science, Slovak University of Agriculture in Nitra, Slovak Republic

ABSTRACT

This study provides information about the method of isolation as well as evaluation of morphology and viability of chicken blastodermal cells (BCs) of Oravka breed. Oravka breed belongs to a critical endangered breed under the conditions of the Slovak Republic. Due to this fact, it is necessary to preserve genetic material of Oravka breed. Frozen blastodermal cells are the potential tool for long-term storage. Chicken blastodermal cells were derived from X stage of embryos (blastoderm stage) using filter paper ring method. To detect blastodermal cells viability specific nuclear fluorochrome Yo-Pro-1 in combination with propidium iodide and DAPI were used. Under the light microscope the isolated blastodermal cells were seen as a large round cells with diameter from 13 to 18 µm and acentric nucleus of 9 µm in diameter. Fluorescent analysis revealed only viable, DAPI-positive cells in the samples. Totally, 38000 blastodermal cells of Oravka breed from each blastoderm were obtained.

Key words: Oravka breed; blastodermal cells; cryopreservation; viability

INTRODUCTION

Conservation of avian genetic resources in the gene bank is necessary for the preservation of endangered species. Cryopreserved biological material can be used for recovering of the lost variation within breeds and restoring of the breeds which have become endangered as a result of destruction of their natural conditions (Sawicka *et al.*, 2011).

Domestic chicken (*Gallus gallus domesticus*) has become an important subject of biotechnological research, particularly in relation to cryopreservation. Compared to mammals, avians have advantages, such as short reproductive cycle, small gap between generations, high laying performance and easy breeding (Li and Lu, 2010). In regards of this facts, avian embryos are

a powerful model to study developmental and stem cell biology.

One of the wonderful advantages of the avian embryo is its accessibility, since the avian embryo is subjected to genetic manipulation (Ishii *et al.*, 2004). Free access to bird eggs (Ishii *et al.*, 2004) allows subjected birds to genetic manipulation Cyriac *et al.*, 2012). This feature has been utilized to produce somatic chimeras of various types through tissue grafts, chorioallantoic grafts, parabiosis, yolk sac chimeras and neural tube chimeras (Le Douarin and McLaren, 1984). However, in order to utilize any potential of avian embryonic stem line for manipulation of the avian genome, it was necessary to develop reliable means of producing germ line chimeras analogous to that produced by blastocyst injection of murine embryos.

***Correspondence:** E-mail: andrea.svoradova@ukf.sk
 Andrea Svoradová, Faculty of Natural Sciences
 Constantine the Philosopher University
 Tr. A. Hlinku 1, 949 74 Nitra, Slovak Republic

Received: February 16, 2016
 Accepted: May 20, 2016

National Agricultural and Food Centre, Research Institute for Animal Production Nitra (NAFC, RIAP) deals with the genetic manipulations of Oravka the local poultry breed. Oravka breed was created by crossbreeding of the local hens in the Orava region with Rhode Island, Wyandotte and New Hampshire breeds (Hanusová *et al.*, 2014) and was recognized in 1990 (Chmelničná, 2004). Weis *et al.* (2010) categorized Oravka breed as a critical endangered breed under the conditions of the Slovak Republic.

The objective of this preliminary study was to optimize the method for blastodermal cells (BCs) isolation in Oravka breed and to evaluate their morphology and viability.

MATERIAL AND METHODS

In our study, chicken eggs of Oravka breed were used as a biological material. X stage blastoderms were collected from 20 freshly laid non-incubated fertilized eggs. The eggs were collected immediately after laying and transferred to the laboratory.

X stage blastoderm was prepared with filter paper ring, the perivitelline membrane was cut around the paper and the blastoderm was removed (Petitte *et al.* 1990). Blastoderm was isolated from yolk and washed several times in calcium-magnesium-free PBS (CMF-PBS) (Applichem, Darmstadt, Germany) to remove as much yolk as possible. Then, the cells were washed and centrifuged in CMF-PBS for 3 min at 300 x g, and supernatant was discarded. The cell concentration was determined using automated cell counter (Z1 Automated Cell Counter, Beckman-Coulter, Germany).

To determine the proportion of viable, apoptotic and dead cells fluorescent staining was used. The total number of cells was determined using 4', 6-diamidino-2-phenylindole (DAPI) fluorescent dye (blue signal; Vector Laboratories, Burlingame, CA, USA). The number of apoptotic cells was determined using Yo-Pro-1 (green signal; Molecular Probes, Lucerne, Switzerland) and the proportion of dead cells using propidium iodide (PI) staining (red signal; Molecular Probes, Eugene, Oregon, USA). Fluorescent dyes were prepared as follows: 1 µl Yo-Pro-1 (1mmol/l) was diluted with 5 µl PI and 194 µl CMF-PBS; 3 µl of cell suspension was transferred to a tube by gentle pipetting and stained with 3 µl Yo-Pro-1/PI by incubation in the dark at room temperature for 20 min. After incubation, sample was washed two times in CMF-PBS. Three µl of the stained sample were placed on a microslide into 3 µl of the Vectashield anti-fade medium containing DAPI fluorescent dye and observed under a LEICA DMI 6000 B fluorescent microscope equipped with DCF 345 FX camera.

RESULTS AND DISCUSSION

Stem cell biology is a rapidly expanding field that includes also the manipulation with stem cells. Chick stem cells can be obtained from embryos and maintained in culture. They can be derived from different sources at various stages of embryonic development. It was demonstrated that in regards to their pluripotency they can form embryoid bodies, which can differentiate into all cell types including embryonic germ layers and contribute to somatic and germline lineages in chimaeras. They are, therefore, comparable to mammalian stem cells, offering a model for studying stem cell biology, as well as being a tool for many applications.

Proper techniques of cell isolation, culture and freezing will give a broad perspective for unlimited storage of genetic recourses.

In our study we isolated, cultured and evaluated under the light microscope BCs from freshly laid non-incubated fertilized eggs of Oravka breed. For isolation we used the same filter paper ring method of BCs isolation as Pettite *et al.* (1990) and Hamidu *et al.* (2010). Our preliminary results showed that this method of chicken BCs isolation is proper and can be used in our subsequent experiments.

Morphological characterization of freshly isolated BCs revealed large round cells with diameter from 13 to 18 µm with acentric nucleus 9 µm in diameter. Furthermore, we determined total number of viable blastodermal cells. Blastoderm of Oravka breed consists of 38000 blastodermal cells. Fluorescent microscopy revealed only living, DAPI positive BCs in the samples (blue signal). Figure 1 (a, b) represents morphology and viability of freshly isolated BCs.

However, for the future most important is the preservation of isolated BCs for long-term storage in liquid nitrogen. Published reports on this topic indicate that BCs tolerate deep-freezing and have ability to produce chicken chimeras (Naito *et al.*, 1992; Bednarczyk *et al.*, 2002).

Chimeras can be used to restore breeds threatened with fate to extinction. Similar studies were conducted by Kino *et al.* (1997) and Naito *et al.* (1994), where injection of BCs into recipient embryos allowed obtaining sex chimeras, and through appropriate mating enabled the restoration of the genotype of the donor. According to Carcience *et al.* (1993) and Naito *et al.* (1992) injected donor blastodermal cells can incorporate into recipient embryos and differentiate into the somatic tissues. The same method was used by Bednarczyk *et al.* (2000); blastoderm was prepared with paper filter rings and thereafter BCs were injected into the sub-germinal cavity of recipient embryo to create transgenic chickens.




Fig. 1: Morphology and viability of freshly isolated BCs. a) light microscope; b) fluorescent microscope – DAPI staining (magnification x 20)

CONCLUSION

Our preliminary study represent only *in vitro* experiments. However, the evaluation of the BCs viability after cryoconservation should be conducted also *in vivo*. The application of isolated BCs for cryopreservation requires further development to expand its potential for avian genetic resources.

ACKNOWLEDGEMENT

This study was financially supported by the Slovak Research and Development Agency (No. APVV-14-0043), VEGA 1/0611/15. This work was supported also by the Research Center AgroBioTech built in accordance with the project Building Research Centre „AgroBioTech“ ITMS 26220220180.

REFERENCES

- BEDNARCZYK, M. – LAKOTA, P. – SIWEK, M. 2000. Improvement of Hatchability of Chicken Eggs Injected by Blastoderm Cells. *Poultry Science*, vol. 79, 2000, p. 1823–1828.
- BEDNARCZYK, M. – LAKOTA, P. – SLOMSKI, R. – PLAWSKI, A. – LIPINSKI, D. – SIEMIENIANKO, B. – CZEKALSKI, P. – GRAJEWSKI, B. – DLUZNIEWSKA, P. 2002. Reconstitution of a chicken breed by inter se mating of germline chimeric birds. *Poultry Science*, vol. 81, 2002, p. 1347–1353.
- CARCIENCE, R. S. – CLARK, M. E. – VERRINDER GIBBINS, A. M. – ETCHES, R. J. 1993. Germline chimeric chickens from dispersed donor blastodermal cells and compromised recipient embryos. *Development*, vol. 117, 1993, p. 669–675.
- CYRIAC, S. – CHURCHIL, R. – R. GIBIN, G. 2012. Transgenic chicken: methods and applications. *Journal of Indian Veterinary Association*, vol. 10 (2), 2012, p. 67–71.
- HAMBURGER, V. – HAMILTON, H. L. 1992. A series of normal stages in the development of the chick embryo. 1951. *Developmental Dynamics*, vol. 195, 1992, p. 231–272.
- HAMIDU, J. A. – RIEGER A. M. – FASENKO G. M. – BARREDA D. R. 2010. Dissociation of chicken blastoderm for examination of apoptosis and necrosis by flow cytometry. *Poultry Science*, vol. 89, 2010, p. 901–909.
- HANUSOVÁ, E. – HRNČÁR, C. – ORAVCOVÁ, M. – HANUS, A. 2014. Characterization of genetic resource in chicken of Oravka breed. *Slovak Journal of Animal Science*, vol. 47, (1), 2014, p. 1–5.
- CHMELNIČNÁ, Ľ. 2004. Oravka. In: KADLEČÍK, O. – HALO, M. – CHMELNIČNÁ, Ľ. – WEIS, J. – KASARDA, R. – MINDEK, S. – MARGETÍN, M. – BULLOVÁ, M. – KOPERNICKÝ, M. – CHLEBO, R. – HRNČÁR, C. SVETLIK, I. – HUBKA, M. 2004: Ohrozené plemená zvierat na Slovensku. Nitra: SPU, 2004, p. 37–52. ISBN 80-8069-459-1.
- ISHII, Y. – REESE, D. E. – MIKAWA, T. 2004. Somatic transgenesis using retroviral vectors in the chicken embryo. *Developmental Dynamics*, vol. 229, 2004, p. 630–642.

- KINO, K. – PAIN, B. – LEIBO, S. B. – COCHRAN, M. – CLARK, M. E. – ETCHES, R. J. 1997. Production of chicken chimeras from injection of frozen-thawed blastodermal cells. *Poultry Science*, vol. 76, 1997, p. 753–760.
- LI, J. – LU, L. 2010. Recent progress on Technologies and applications of transgenic poultry. *African Journal of Biotechnology*, vol. 9 (24), 2010, p. 3481–3488.
- NAITO, M. – NIRASAWA, K. – OISHI T. 1992. Preservation of quail blastoderm cells in liquid nitrogen. *British Poultry Science*, vol. 33, 1992, p. 449–453.
- NAITO, M. – TAJIMA A. – TAGAMI, T. – YASUDA, Y. – KUWANA, T. 1994. Preservation of chick primordial germ cells in liquid nitrogen and subsequent production of viable offspring. *Journal of Reproduction and Fertility*, vol. (102), 1994, p. 321–325.
- PETITTE, J. N. 1990. Production of somatic and germline chimeras in the chicken by transfer of early blastodermal cells. *Development*, vol. 108, 1990, p. 185–189.
- SAWICKA, D. – BRZEZIŃSKA, J. – BEDNARCZYK, M. 2011. Cryoconservation of embryonic stem cells and gametes as a poultry biodiversity preservation method. *Folia Biologica*, vol. 5, 2011, p. 1–5.
- WEIS, J. – GARDIÁNOVÁ, I. – HRNČÁR, C. – MINDEK, S. – SVOBODOVÁ, I. – BUJK, O. J. 2010. Analýza početnosti autochtónnych plemien kúr na území Slovenskej republiky. *Acta fytotechnica et zootechnica*, roč. 10, 2010, s. 31–33.

Slovak Journal of Animal Science

<http://www.vuzv.sk/index.php/slovak-journal-of-animal-science>

<http://www.sjas.sk>

INSTRUCTIONS FOR AUTHORS

The scientific journal Slovak Journal of Animal Science publishes original papers, review articles, short communications, reviews of important works, chronicles, and reports on participation in important international conferences. Topics of the journal include problems in biology of animals, animal production and veterinary medicine.

The author is responsible for originality of the paper, subject-matter and formal attributes of the paper. Papers are published in English language. The author is responsible for translation into English. In case of linguistic shortcomings the paper will be sent back to the author.

Copyright. The journal is protected by copyright held by the publisher after the manuscript has been accepted for publication. As regards the transfer of rights, the corresponding author assumes responsibility for all the authors.

MANUSCRIPT SUBMISSION

- **Manuscripts** should be sent by e-mail as attachments. Alternatively, they can be submitted on floppy disk, USB key, in hard copy and a properly labeled Compact Disk (CD) with identical contents, including figures. Unique MS Word file must be saved under the first author's surname only. They are to be sent to the following address:

Slovak Journal of Animal Science - Editorial Office
National Agricultural and Food Centre - Research Institute
for Animal Production Nitra
Hlohovecká 2, 951 41 Lužianky
Slovak Republic

Phone: +421 37 6546 249 Fax: +421 37 6546 311 e-mail: editor@vuzv.sk

- **The author's declaration** (find at www.vuzv.sk/index.php/slovak-journal-of-animal-science) that the paper has not been published anywhere else should be enclosed. The Declaration must be carefully completed and signed by the first author.

- **The Corresponding Author** should include his or her full name including all academic, scientific and pedagogic titles and the detailed address of the institution with postal code, telephone and fax numbers, and e-mail address.

REVIEWERS' EVALUATION

All scientific contributions will be peer-reviewed on the criteria of originality and quality. Two reviewers, who have appropriate knowledge of the subject, are appointed for each paper. If need be third referee will be asked. The reviewers are asked to return papers within 3 weeks. Editorial board of the journal decides on contributing publications, taking into consideration the standpoint of reviewers, scientific significance and quality of manuscript. The manuscript can be rejected if it does not comply with the subject of the journal. The editor makes every effort to explain to authors the reasons of rejection.

AFTER REVIEWERS' EVALUATION

After the author receives the reviewers' opinion he owes to return the corrected manuscript to the editorial office within three weeks. The authors may, however, request an extension of the re-submission deadline if necessary. After incorporation of revisions it is necessary to send the new complete manuscript with tables and figures, also in case of minor revisions in the original manuscript. In an enclosure the author has to answer all the important comments of the reviewers clearly and in detail. The author is not obliged to respect the comments of reviewers but he has to explain in the enclosure why he does not accept the reviewers' comments. If the author does not keep the term of sending back the revised manuscript, his paper is liable to be excluded from further processing.

SENDING THE CORRECTED PROOF

The author has to return the made-up manuscript for proof-reading. He must send it back within two days. Within this stage of manuscript preparation for printing, corrections are done only for such errors that arise during the work in editorial office.

FORMAL LAYOUT OF PAPER

TITLE

The title should briefly but accurately characterize the aim of the paper. It is not advisable to use any abbreviation in the title.

AUTHOR(s)

Names of authors should be without titles, in capital letters; first name at first and then surname. Individual authors separated by commas. *E.g.* J. RAFAY, K. BEZOVA, P. FLAK

ABSTRACT – subtitle

They should contain abbreviated exact subject of the paper without any added interpretation or criticism. It must (together with tables and figures that are published in the text, possibly with a reference to them) give the reader sufficient amount of information about the paper, namely about the objective, method and the most important results of the work (mentioned are statistical and numeric results). The abstract must not exceed 350 words.

KEY WORDS

They are arranged from more general terms towards more precise ones. They begin with small letter and are separated by semicolon. They should contain from 3 to 6 words.

INTRODUCTION – subtitle

Shall give brief and clear information about the current state of the research in the given field in our country and abroad (references of authors) and about the objective of the work with explanation of scientific or social necessity to solve it. Quotation should serve as connection of the quoted place with the References. In the scientific journal Slovak J.Anim.Sci. the “method of utilization of the first entry and date” is used for quotation in agreement with the norms. Names of authors are written with small letters in brackets (*e.g.* Novak, 1999). It is recommended to include references to papers from peer periodicals only. Citations from nonavailable sources (reports, national journals, proceedings, etc.) should be omitted. Papers published by one or two authors are to be cited by their names, those published by three or more authors by the name of the first one, *et al.* If more than one paper by the same author/two authors/first author, *et al.*, published in the same year is cited, they should be differentiated by YEAR_{a,b,c} both in the text and the list of References. Citation in text: *e.g.*: Clark and Davis (1980), Citation within brackets: In case of more than one cited author in one paper, only first author and *et al.* are referred, *e.g.*: (Tesařík *et al.*, 1986) If the same problem is solved by several authors in separate works, then they are separated by semicolon, *e.g.*: (Blaho *et al.*, 1980; Davis, 1990; Hyttel *et al.*, 2000;) Several papers cited together should be arranged according to the year of publication starting with the oldest one.

MATERIAL AND METHODS – subtitle

Should describe the studied material as precisely as possible, as well as used methods of study and ways of evaluation of results. Used method of experiment is described only in case of original or modified method, or with quotation of author of the method. The description of method should enable the interested specialist to repeat the work. Statistical methods must be described in details including the software used.

RESULTS – subtitle

In this part are mentioned only concrete results of work, and in text are mentioned also the most important numerical values and data mentioned in tables. Only significant differences should be commented. It is necessary to support the results by variation- statistical processing. Statistical evaluation of results must be logically explained and perfectly processed. Results should contain data that make it possible to verify them (*e.g.* with evaluation of significance of differences in two averages, it is necessary to give the average, number of determinations and standard deviation). Statistically significant differences must be unambiguously and clearly marked in tables or graphs.

DISCUSSION – subtitle

It is necessary to assess results from various aspects mainly with respect to factors that could influence them, and to confront them with results of other authors in such a way as to show what is new in the obtained results and what differs them from results of other authors or hitherto knowledge or in which part they correspond to each other. Bibliographical reference of authors should be in the same form as in the introductory part.

RESULTS AND DISCUSSION May comprise of a chapter

CONCLUSION – subtitle

It has to be clear and brief. It should define clearly the obtained new knowledge, the most important results, their scientific or practical significance and recommendations for the practice or for further research.

ACKNOWLEDGEMENT

At the end of the paper should be mentioned the number and title of project within which the problem was studied and the published results obtained.

REFERENCES – subtitle

Bibliographical reference – set of data enabling identification of document that is connected in some way with creation of another document. References should be in alphabetical order after the surname of the first author. The references are listed after the title of the paper if there is no name of an author. Listed are only sources that are quoted in the text. Selected literary sources should be directly connected to the problems dealt with in the paper and they should consist only of peer-reviewed scientific journals. Listed should be only primary sources, based on results of the quoted authors. Bibliographic references and quotations shall be written according to the international standards STN ISO 690. References in the text should agree with those in the list of references.

Editorial board of the journal set the following rules:

- in References are listed names of all authors;
- in papers are used following abbreviations: *et al.*; 2nd ed., 3rd ed., ...6th ed...; vol., no., p.

EXAMPLES OF REFERENCES:

Example: Monograph

RICKE, S. C. – VAN LOO, E. J. – JOHNSON, M. G. – O'BRYAN, C. A. 2012. *Organic Meat Production and Processing*. The 1st edition. USA: Wiley-Blackwell, 2012, 464 p. ISBN 978-0-8138-2126-9.

Example: Paper in monograph

PRICE, D. 1980. A general theory of bibliometric and other cumulative advantage processes. In: GRIFFITH, B.: *Key papers in information science*. New York : Knowledge Industry Publications, 1980, p. 177–191.

Example: Paper in journal

BROUCEK, J. – UHRINCAT, M. – TANCIN, V. – HANUS, A. – TONGEL, P. – BOTTO, L. – BOZIK, I. 2013b. Performance and behaviour at milking after relocation and housing change of dairy cows. *Czech Journal of Animal Science*, vol. 58, 2013b, p. 389–395.

Example: Article in collected papers

KUŽELOVÁ, L. – CHRENEK, P. 2014. Effect of rooster macrophages on the motility characteristics of spermatozoa. In: *International Symposium on Animal Science* : Proc. Belgrade : University of Belgrade, Serbia, 2014, p. 73-78. ISBN 978-86-7834-199-1.

Compliance with these instructions is obligatory for all authors.

If a manuscript does not comply with the above requirements, the editorial office will not accept it for consideration and will return it to the authors.

Slovak Journal of Animal Science
<http://www.vuzv.sk/index.php/slovak-journal-of-animal-science>
<http://www.sjas.sk>

Subscription and distribution

All orders should be sent direct to:
Slovak Journal of Animal Science
Editorial Office
NPPC - Research Institute for Animal
Production Nitra
Hlohovecká 2
951 41 Lužianky, Slovak Republic
Phone: +421 37 6546 249
Fax: +421 37 6546 311
E-mail: editor@vuzv.sk

Subscription rates 2016 (journal + postage)

Print:

65,- € collecting in editorial office directly
75,- € Slovak Republic
85,- € Czech Republic
95,- € Europe
105,- € ROW

Offprints:

Editorial office will send on request the offprint
in Portable Document Format (PDF) via e-mail.

Contents

Original papers

- OLEXIKOVÁ, L. – ŠPALEKOVÁ, E. – KUBOVIČOVÁ, E. – MAKAREVICH, A. V. – CHRENEK, P.: 57
Fertilizing ability of Pinzgau bull sperm *in vitro* after cryostorage
- KOVÁČ, M. – KULÍKOVÁ, B. – VAŠÍČEK, J. – CHRENEK, P.: 62
Cryopreservation of amniotic fluid stem cells derived from Zobor rabbit
- LALA, A. O. – OKWELUM, N. – BELLO, K. O. – FAMAKINDE, N. A. – ALAMU, M. O.: 68
Comparative study between Isa Brown and Fulani ecotype chickens supplemented with humic acid
- OJO, V. O. A. – OYEBANJO, E. D. – OPAYEMI, O. T. – MUSTAPHA, S. O. – ADELUSI, O. O. 76
– ADEOYE, S. A. – ANELE, U. Y. – OGUNSAKIN, A. O. – JOLAOSHO, A. O. – ONIFADE, O. S.:
Effects of seasonal changes on the nutritive quality of *Moringa oleifera* and *Panicum maximum* silage

Review

- TANČIN, V. – BAUER, M. – HOLKO, I. – BARANOVIČ, Š.: 85
Etiology of mastitis in ewes and possible genetic and epigenetic factors involved

Short communication

- SVORADOVÁ, A. – KUŽELOVÁ, L. – CHRENEK, P.: 94
The potential use of chicken blastodermal cells of Oravka breed